


Please type a plus (+) inside this box - 

PTO/SB/29 (12/97)

Approved for use through 09/30/00, OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

09/22/99
JCE44 U.S. PRO

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 1803-247 Total Pages 160

First Named Inventor or Application Identifier

MAURICE K. GATELY

Express Mail Label No. EL168272336US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

1. ☒ Fee Transmittal Form
Submit an original, and a duplicate for fee processing)
2. ☒ Specification [Total Pages 105]

(preferred arrangement set forth below)

- Descriptive title of the Invention
- Cross Reference to Related Applications
- Statement Regarding Fed sponsored R&D
- Reference to Microfiche Appendix
- Background of the Invention
- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description of the Invention (including drawings, if filed)
- Claim(s)
- Abstract of the Disclosure

3. ☒ Drawing(s) (35 USC 113) [Total Sheets 44]

4. ☒ Oath or Declaration [Total Sheets 04]

- a. ☐ Newly executed (original or copy)
b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for divisional with Box 17 completed)

[Note Box 5 below]

- i. ☐ DELETION OF INVENTOR(S)

Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33 (b).

5. ☒ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. ☐ Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)

- a. ☐ Computer Readable Copy
b. ☐ Paper Copy (identical to computer copy)
c. ☐ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (copy from prior application)
9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney (when there is an assignee)
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
12. ☒ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
14. ☐ Small Entity ☐ Statement filed in prior application, Statement(s) Status still proper and desired
15. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
16. ☒ Other: Copy of Petition For Extension of Time from parent 08/459,151 application; copy of associate Power of Attorney from 08/459,151 application.

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☒ Divisional ☐ Continuation-in-part (CIP) of prior application No: 08/459,151, filed June 2, 1995

18. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

20583

(Insert Customer No. or Attach bar code label here)

or ☐ Correspondence address below

NAME			
ADDRESS			
CITY	STATE	ZIP CODE	
COUNTRY	TELEPHONE	FAX	

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Prior application: Examiner P. Mertz, Ph.D.Art Unit 1646

Assistant Commissioner for Patents
 Box PATENT APPLICATION
 Washington, D.C. 20231

Sir:

This is a request for filing a ☐ continuation ☒ divisional application under 37 CFR § 1.53(b), of pending prior application no. 459,151, filed on June 2, 1995, which is a divisional of application Serial No. 205,011, filed March 2, 1994, now abandoned, which is a divisional of application Serial No. 857,023, filed March 24, 1992, now abandoned, which is a continuation-in-part of application Serial No. 572,284, now abandoned, which is a continuation-in-part of application Serial No. 520,935, filed May 9, 1990, now abandoned, which is a continuation-in-part of application Serial No. 455,708, filed December 22, 1989, now abandoned, each of which is incorporated herein by reference in its entirety.

of Maurice K. Gately, Ulrich A. Gubler, Jeffrey D. Hulmes, Frank J. Podlaski, Alvin S. Stern, Richard A. Chi and Yu-Ching E. Pan

(inventor(s) currently of record in prior application)

for PURIFICATION AND CHARACTERIZATION OF CYTOTOXIC LYMPHOCYTE
MATURATION FACTOR AND MONOCLONAL ANTIBODIES THERETO

(title of invention)

1. ☒ The filing fee is calculated below, and reflects the pending claims as amended by the accompanying Preliminary Amendment:

PATENT APPLICATION FEE VALUE


TYPE	NO. FILED	LESS	EXTRA	EXTRA RATE	FEE
Total Claims	6	-20	0	\$18.00 each	0.00
Independent	3	-3	0	\$78.00 each	0.00
Basic Fee					760.00
Multiple Dependency Fee If Applicable (\$270.00)					
Total					0.00
50% Reduction for Independent Inventor, Nonprofit Organization or Small Business Concern					-
Total Filing Fee					\$ 760.00

2. ☒ Please charge the required fee to Pennie & Edmonds LLP Deposit Account No. 16-1150. A copy of this sheet is enclosed.
3. ☐ Amend the specification by inserting before the first line the following sentence:
- 4a. ☐ Transfer the drawings from the prior application to this application and abandon the prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file.
- 4b. ☐ New formal drawings are enclosed.
- 4c. ☒ Informal drawings are enclosed.
- 5a. ☐ Priority of application no. filed on is claimed under 35 U.S.C. §119.
- 5b. ☐ The certified copy has been filed in prior application no. , filed .
6. ☐ The prior application is assigned of record to.
- 7a. ☒ A copy of the Power of Attorney appears in the original papers in the prior application no. 08/459,151, filed June 2, 1995. A copy of the executed Power of Attorney is being submitted herewith.
- 7b. ☐ Since the Power of Attorney does not appear in the original papers, a copy of the Power in prior application no. , filed is enclosed.
8. ☐ This application contains nucleic acid and/or amino acid sequences required to be disclosed in a Sequence Listing under 37 CFR §§1.821-1.825. It is requested that the Sequence Listing in computer readable form from prior application no. , filed , be made a part of the present application as provided for by 37 C.F.R. §1.821(e). The sequences disclosed therein are the same as the sequences disclosed in this application. A copy of the paper Sequence Listing from application no. is enclosed.
9. ☐ The undersigned states, under 37 C.F.R. §1.821(f), that the content of the enclosed paper Sequence Listing from application no. is the same as the content of the computer readable form submitted on in application no. .

10. ☒ Additional enclosures or instructions:
- A) Preliminary Amendment;
 - B) Copy of a Petition for Extension of Time filed in connection with parent application 08/459,151 requesting that the period for response to a pending Office Action be extended from August 12, 1999 up to and including November 12, 1999;
 - C) A Request Under 37 C.F.R. §§ 1.607 and 1.608(a) for Interference With a Patent; and
 - D) A copy of an Associate Power of Attorney filed in connection with parent application 08/459,15.

Respectfully submitted,

September 22, 1999


Thomas E. Friebe 29,258
(Reg. No)
PENNIE & EDMONDS LLP
1155 Avenue of the Americas
New York, NY 10036

Tel. No. (212) 790-9090

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: GATELY, MAURICE K. et al.

Serial No.: To Be Assigned
(Division Of Serial No.
08/459,151)

Group Art Unit: To Be
Assigned

Examiner: To Be Assigned

Filed: On Even Date Herewith

Attorney Docket No.: 1803-247

For: PURIFICATION AND
CHARACTERIZATION OF
CYTOTOXIC LYMPHOCYTE
MATURATION FACTOR AND
MONOCLONAL ANTIBODIES
THERETO

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Pursuant to the Rules of Practice, please enter the following amendments and remarks into the file of the above-captioned application, which is being filed concurrently herewith.

Applicants also submit herewith a Request Under 37C.F.R. §§ 1.607 and 1.608(a), and its accompanying documents.

IN THE SPECIFICATION:

Please amend the specification as follows:

Delete the first sentence following the title ("This is a Continuation-In-Part...") and insert in its place the following sentence:

--This is a Divisional of application Serial No. 459,151, filed June 2, 1995, which is a Divisional of application Serial No. 205,011, filed March 2, 1994, now abandoned, which is a Divisional of application Serial No. 857,023, filed March 24, 1992, now abandoned, which is a Continuation-In-Part of application Serial No. 572,284, now abandoned, which is a Continuation-In-Part of application Serial No. 520,935, filed May 9, 1990, now abandoned, which is a continuation-in-part of application Serial No. 455,708, filed December 22, 1989, now abandoned, the contents of each of which is incorporated herein by reference in its entirety.-

On page 9, line 28, delete "25" and insert in its place --25A-25D--.

On page 9, line 31, delete "26" and insert in its place --26A-26C--.

On page 12, line 30, delete "23 and 24" and insert in its place --25A-25D and 26A-26C--.

On page 64, line 11, delete "25" and insert in its place --25A-25D--.

On page 67, line 7, delete "26" and insert in its place --26A-26C--.

IN THE CLAIMS:

Please amend the claims as follows:

Cancel Claims 1-32, without prejudice.

Add new Claims 33-38, as follows:

--33. A monoclonal antibody which specifically reacts with cytotoxic lymphocyte maturation factor (CLMF) protein, said protein comprising:

- (a) a first subunit having an apparent molecular weight of approximately 40 kD under reducing conditions on SDS PAGE and comprising the amino acid sequence of FIG. 25A-25D from amino acids 23 to 328; and
- (b) a second subunit having an apparent molecular weight of approximately 30-35 kD under reducing conditions on SDS PAGE and comprising the amino acid sequence of FIG. 26A-26C from amino acid 23 to 219.

34. The monoclonal antibody of Claim 33 wherein the CLMF protein is capable of inducing proliferation of phytohemagglutinin (PHA)-activated peripheral blood lymphocytes.

35. The monoclonal antibody of Claim 33 wherein said antibody reacts with said first subunit.

36. The monoclonal antibody of Claim 33 wherein said antibody reacts with said second subunit.

37. A monoclonal antibody which specifically reacts with a subunit of cytotoxic lymphocyte maturation factor (CLMF) protein, said subunit having an apparent molecular weight of approximately 40 kD under reducing conditions on SDS PAGE and comprising the amino acid sequence of FIG. 25 from amino acids 23 to 328.

38. A monoclonal antibody which specifically reacts with a subunit of cytotoxic lymphocyte maturation factor (CLMF) protein, said subunit having an apparent molecular weight of approximately 30-35 kD under reducing conditions on SDS PAGE and comprising the amino acid sequence of FIG. 26 from amino acid 23 to 219.--

REMARKS

This application is a Divisional of application Serial No. 459,151 (the "'151 application"), filed June 2, 1995, which is a Divisional of application Serial No. 205,011, filed March 2, 1994, now abandoned, which is a Divisional of application Serial No. 857,023, filed March 24, 1992, now

abandoned, which is a Continuation-In-Part of application Serial No. 572,284, now abandoned, which is a Continuation-In-Part of application Serial No. 520,935, filed May 9, 1990, now abandoned, which is a continuation-in-part of application Serial No. 455,708, filed December 22, 1989, now abandoned.

Applicants note that a Petition for Extension of Time has been timely filed in the '151 application, extending time for response to an Office Action pending in the '151 application for three months, from August 12, 1999 up to and including November 12, 1999.

With this Preliminary Amendment, Claims 1-32, which were pending in the parent '151 application, have been canceled without prejudice to Applicants' right to pursue the subject matter of the canceled claims in other applications, and have been replaced with new Claims 33-38. Thus, Claims 33-38 are currently pending in this application.

Claims 33-38 are directed to monoclonal antibodies which specifically react with cytotoxic lymphocyte maturation factor (CLMF). Claims 33-38 are fully supported by the instant specification.

Claims 33 and 34 are directed to a monoclonal antibody which specifically reacts with CLMF protein, said CLMF protein comprising a subunit of 40 kD under reducing conditions comprising the amino acid sequence of instant application Figure 25A-25D from amino acid residues 23 to 328, and a subunit of 30-35 kD under reducing conditions comprising

the amino acid sequence of instant application Figure 26A-26C from amino acid residues 23 to 219. Claim 34, which depends from Claim 33, recites one of the biological activities of the CLMF protein, in particular that the CLMF protein can induce proliferation of phytohemagglutinin (PHA)-activated peripheral blood cells.

These claims are fully supported by the instant application. Working Example 13 (pp. 71-78) describes the successful generation and characterization of monoclonal antibodies directed against CLMF. The structural features and biological activity of the CLMF protein recited in the claims are as characterized and described in the 'instant application. For a characterization of the size and amino acid sequence of the two CLMF subunits, see, e.g., p. 33, ll. 10-12 and Figure 7; p. 64, ll. 10-17 and Fig. 25; and p. 67, ll. 5-15 and Fig. 26. The ability of the CLMF protein to induce proliferation of PHA-activated peripheral blood cells (referred to in the specification as a "T cell growth factor (TGF) assay") is described and demonstrated at p. 19, l. 15 to p. 21, l. 2, and in working Example 9 (pp. 54-60).

Claim 35, which depends from Claim 33, and independent Claim 37 are directed to monoclonal antibodies that react with the 40 kD CLMF subunit. Working example 13 describes the production and characterization of monoclonal antibodies which specifically react with the 40 kD CLMF subunit. See especially p. 73, l. 20 to p. 74, l. 9.


Claim 36, which depends from Claim 33, and independent Claim 37 are directed to monoclonal antibodies that react with the 35 kD CLMF subunit. Example 14 (pp. 79-80) describes a method for producing monoclonal antibodies which specifically react with the 35 kD CLMF subunit, wherein the monoclonal antibodies are generated against a synthetic peptide containing a 35 kD CLMF subunit amino acid sequence.

The specification has been amended to update the related application information and to adjust figure legend designations accordingly.

The amendments made herein do not constitute new matter. Applicants respectfully request that the amendments and remarks made herein be entered into the record of the above-captioned patent application.

Respectfully submitted,

Date: September 22, 1999


Thomas E. Friebe 29,258
(Reg. No.)

PENNIE & EDMONDS
1155 Avenue of the Americas
New York, New York 10036-2711
(212) 790-9090

PURIFICATION AND CHARACTERIZATION OF
CYTOTOXIC LYMPHOCYTE MATURATION FACTOR
AND MONOCLONAL ANTIBODIES THERETO

This is a Continuation-in-Part of Serial Number 07/572,284,
filed August 27, 1990, which is a continuation-part of Serial No.
10 07/520,935, filed May 9, 1990, which is a Continuation-in-Part of
Serial Number 07/455,708, filed December 22, 1989.

FIELD OF THE INVENTION

15 The present invention relates to the field of cytokines, in
particular to those cytokines which synergize with Interleukin-2 (IL-
2) to activate cytotoxic lymphocytes.

BACKGROUND OF THE INVENTION

20 'Cytokine' is one term for a group of protein cell regulators,
variously called lymphokines, monokines, interleukins and
interferons, which are produced by a wide variety of cells in the
body, play an important role in many physiological responses, are
25 involved in the pathophysiology of a range of diseases, and have
therapeutic potential. This heterogeneous group of proteins has the
following characteristics in common. They are low molecular weight
(≤ 80 kDa) secreted proteins which are often glycosylated; are
involved in immunity and inflammation where they regulate the
30 amplitude and duration of a response; and are usually produced
transiently and locally, acting in a paracrine or autocrine, rather than
endocrine manner. Cytokines are extremely potent, generally acting
at picomolar concentrations; and interact with high affinity cell
surface receptors specific for each cytokine or cytokine group. Their
35 cell surface binding ultimately leads to a change in the pattern of

cellular RNA and protein synthesis, and to altered cell behavior. Individual cytokines have multiple overlapping cell regulatory actions.

- 5 The response of a cell to a given cytokine is dependent upon the local concentration of the cytokine, the cell type and other cell regulators to which it is concomitantly exposed. The overlapping regulatory actions of these structurally unrelated proteins binding to different cell surface receptors is at least partially accounted for by
- 10 the induction of common proteins which can have common response elements in their DNA. Cytokines interact in a network by: first, inducing each other; second, transmodulating cytokine cell surface receptors and third, by synergistic, additive or antagonistic interactions on cell function. [Immunology Today 10 No. 9 p 299
- 15 (1989)].

- The potential utility of cytokines in the treatment of neoplasia and as immunoenhancing agents has recently been demonstrated in studies using human recombinant interleukin-2 (rIL-2). Natural
- 20 Interleukin-2 (IL-2) is a lymphokine which is produced and secreted by T-lymphocytes. This glycoprotein molecule is intimately involved in the induction of virtually all immune responses in which T-cells play a role. B cell responses *in vitro* are also enhanced by the presence of IL-2, and IL-2 has also been implicated as a
- 25 differentiation inducing factor in the control of B and T lymphocyte responses.

- Administration of human rIL-2 has been shown in some cases to result in regression of established tumors in both experimental
- 30 animals [J. Exp. Med 161:1169-1188, (1985)] and in man [N. Engl. J. Med 313: 1485-1492, (1985) and N. Engl. J. Med 316:889-897 (1987)]. The anti-tumor effects of rIL-2 are thought to be mediated by host cytotoxic effector lymphocytes which are activated by rIL-2
- 35 *in vivo* [J. Immunol. 139:285-294 (1987)]. In addition, results from animal models suggest that rIL-2 might also have value in the treatment of certain infectious diseases [J. Immunol. 135:4160-4163

(1985) and J. Virol. 61:2120-2127 (1987)] and in ameliorating chemotherapy-induced immunosuppression [Immunol Lett. 10:307-314 (1985)].

- 5 However, the clinical use of rIL-2 has been complicated by the serious side effects which it may cause [N. Engl. J. Med. 313:1485-1492 (1985) and N. Engl. J. Med. 316:889-897 (1987)]. One approach to improving the efficacy of cytokine therapy while reducing toxicity is to use two or more cytokines in combination. For example,
- 10 synergistic antitumor activity has been shown to result when rIL-2 is administered to tumor-bearing mice together with recombinant interferon alpha (rIFN alpha) [Cancer Res. 48:260-264 (1988) and Cancer Res. 48:5810-5817 (1988)] or with recombinant tumor necrosis factor alpha (rTNF alpha)[Cancer Res. 47:3948-3953 (1987)].
- 15 Since the antitumor effects of IL2 are thought to be mediated by host cytotoxic effector lymphocytes, it would be of interest to identify and isolate novel cytokines which synergize with rIL2 to activate cytotoxic lymphocytes *in vitro*. These novel cytokines would also be useful as antitumor agents when administered in combination with
- 20 rIL2 *in vivo*.

SUMMARY OF THE INVENTION

- The present invention is a novel cytokine protein called
- 25 Cytotoxic Lymphocyte Maturation Factor (CLMF) also called IL-12 which is produced and synthesized by cells capable of secreting CLMF such as mammalian cells particularly human NC-37 B lymphoblastoid cells (ATCC CCL 214 American Type Culture Collection, Rockville, MD). CLMF synergistically induces with low concentrations
- 30 of IL-2 (IL-2) the cytolytic activity of Lymphokine Activated Killer (LAK) cells, and CLMF is capable of stimulating T-cell growth.

 The present invention is directed toward the process of isolating CLMF in a substantially pure form.

The process comprises the following:

stimulating NC-37 B lymphoblastoid cells to produce and secrete cytokines into a supernatant liquid;

5

collecting the supernatant liquid produced by the stimulated cells;

separating the supernatant liquid into protein fractions;

10

testing each protein fraction for the presence of CLMF;

retaining the protein fractions which are able to stimulate T-cell growth, said fractions containing an active protein which is responsible for the T-cell stimulating activity of the protein fractions;

15

isolating said active protein into a substantially pure form, said protein being Cytolytic Lymphocyte Maturation Factor (CLMF). CLMF is a 75 kilodalton (kDa) heterodimer comprised of two polypeptide subunits, a 40 kDa subunit and a 35 kDa subunit which are bonded together via one or more disulfide bonds.

20

The process of this invention is capable of purifying CLMF from any liquid or fluid which contains CLMF together with other proteins. Also claimed are the protein fractions capable of stimulating T-cell growth, the substantially purified active protein, CLMF, obtained from the above described process,

25

the isolated cloned genes encoding the 40 kDa subunit and the 35 kDa subunit, vectors containing these genes and host cells transformed with the vectors containing the genes.

30

In addition a method for stimulating LAK cells and T-cells comprised of treating these cells with CLMF alone or with IL-2 is claimed. Also claimed are isolated antibodies capable of binding to CLMF.

5

Antibodies to CLMF

Monoclonal antibodies prepared against a partially purified preparation of CLMF have been identified and characterized by 1: immunoprecipitation of ^{125}I -labelled CLMF, 2: immunodepletion of CLMF bioactivity, 3: western blotting of CLMF, 4: inhibition of ^{125}I -CLMF binding to its cellular receptor and 5: neutralization of CLMF bioactivity. Twenty hybridomas were identified which secreted anti-CLMF antibodies. The antibodies immunoprecipitate ^{125}I -labelled CLMF and immunodeplete CLMF bioactivity as assessed in the T-cell proliferation and LAK cell induction assays. Western blot analysis demonstrate that each antibody binds to the 70 kDa heterodimer and to the 40 kDa subunit. These data demonstrated that the 20 antibodies were specific for CLMF and in particular for the 40 kDa subunit of CLMF. A CLMF receptor binding assay has been developed to evaluate the ability of individual antibodies to inhibit CLMF binding to its cellular receptor. The assay measures the binding of ^{125}I -labelled CLMF to PHA activated PBL blast cells in the presence and absence of each antibody. Of the 20 antibodies tested, 12 antibodies inhibit greater than 60% of the ^{125}I -labelled CLMF binding to the blast cells. Two inhibitory antibodies, 7B2 and 4A1, neutralize CLMF bioactivity while one non-inhibitory antibody, 8E3, does not neutralize CLMF bioactivity. These data confirm that antibodies which block ^{125}I -labelled CLMF binding to its cellular receptor will neutralize CLMF bioactivity as assessed by the T-cell proliferation and LAK cell induction assays. The ability of the antibodies specific for the 40 kDa subunit of CLMF to neutralize CLMF bioactivity indicates that determinants on the 40 kDa subunit are necessary for binding to the CLMF cellular receptor.

35

Utility for the Monoclonal Anti-Human CLMF Antibodies

- The monoclonal anti-CLMF antibodies provide powerful analytical, diagnostic and therapeutic reagents for the
- 5 immunoaffinity purification of natural and recombinant human CLMF, the development of human CLMF immunoassays, the identification of the active site of the 40 kDa subunit of CLMF and as possible therapeutic treatments which require selective immunosuppression of cytotoxic T cells, such as in transplantation.
- 10 Monoclonal antibodies which recognize different epitopes on human CLMF can be used as reagents in a sensitive two-site immunoassay to measure levels of CLMF in biological fluids, cell culture supernatants and human cell extracts.
- 15 The present invention is directed to monoclonal antibodies against CLMF which exhibit a number of utilities including but not limited to:
1. Utilizing the monoclonal antibodies as affinity reagents for
20 the purification of natural and recombinant human CLMF;
 2. Utilizing the monoclonal antibodies as reagents to configure enzyme-immunoassays and radioimmunoassays to measure natural and recombinant CLMF in biological fluids, cell culture supernatants,
25 cell extracts and on plasma membranes of human cells and as reagents for a drug screening assay;
 3. Utilizing the monoclonal antibodies as reagents to construct sensitive two-site immunoassays to measure CLMF in biological
30 fluids, cell culture supernatants and human cell extracts;
 4. Utilizing the monoclonal antibodies as reagents to identify determinants of the 40 kDa subunit which participate in binding to the 35 kDa subunit and which participate in binding to the CLMF
35 cellular receptor;

5. Utilizing the intact IgG molecules, the Fab fragments or the humanized IgG molecules of the inhibitory monoclonal antibodies as therapeutic drugs for the selective blockade of proliferation and activation of cytotoxic T cells, such as in transplantation.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a plot of a supernatant solution obtained from cultured NC37 lymphoblastoid cells applied to a Nu-Gel P-SP column
10 showing the protein fraction containing TGF activity being eluted with a salt gradient.

Figure 2 is a plot of the material containing TGF activity obtained from the separation of Figure 1 as it was being eluted with
15 a salt gradient through a Blue-B-Agarose Column.

Figure 3 shows the plot of the material containing TGF activity obtained from the separation of Figure 2 as it was being eluted with a NaCl gradient through a Mono Q column.
20

Figure 4 shows the SDS-polyacrylamide gel electrophoresis (SDS-PAGE) analysis of the fractions obtained from the step illustrated in Figure 3.

Figure 5 shows the elution profile through a Vydac Diphenyl column of fraction 38 from the Mono Q Chromatography separation.
25

Figure 6 shows SDS-PAGE analysis of protein purity of the protein fractions 86-90 recovered from the process depicted in
30 Figure 5.

Figure 7 shows the SDS PAGE analysis of fractions 87 and 88 under non-reducing (without β -mercaptoethanol) and reducing (in the presence of β -mercaptoethanol) conditions showing the 75,000
35 molecular weight CLMF separated into two subunits of 40 kDa and 35 kDa.

Figure 8 shows the elution pattern of the proteins from the supernatant solution from NC-37 cells applied to a Nu- Gel P-SP column eluted with a salt gradient in Example 2.

5

Figure 9 is a Blue-B-Agarose column salt gradient elution profile of the active fractions obtained from the Nu-Gel P-SP column elution of Figure 8.

10 Figure 10 is a Mono-Q column salt gradient elution profile of the active fractions obtained from the elution shown in Figure 9.

Figure 11 is the elution pattern through a Vydac Diphenyl column of active fractions 39 and 40 obtained from the Mono Q
15 Chromatography shown in Figure 10.

Figure 12 is the SDS PAGE analysis under reducing conditions of the active fractions obtained from the elution of Figure 11.

20 Figure 13 is a schematic diagram showing the separation of the 40kDa subunit from the 35kDa subunit of the CLMF cytokine of the present invention.

25 Figure 14 is a schematic diagram showing the determination of the amino acid composition, N-terminal sequencing, proteolytic digestion and complete sequencing of the 40kDa subunit of the CLMF cytokine of the present invention.

30 Figure 15 is a tryptic peptide map of the digested 40kDa subunit of the CLMF cytokine of the present invention.

Figure 16 is a proteolytic peptide map of the digested 40kDa subunit CLMF in which the proteolytic enzyme which was used was Staphylococcus aureus V8 protease.

35

Figure 17 is a chart which summarizes the protein structural determination of the 40kDa subunit of CLMF.

Figure 18 is the SDS PAGE analysis of Fraction 39 from the Mono Q FPLC elution shown in Figure 4.

Figure 19 is the elution pattern through a Vydac C-18 column of fraction 39 of the Mono Q chromatography which was reduced in 5% β -mercaptoethanol.

Figure 20 is the SDS-PAGE gel analysis under non-reducing conditions of the fractions which were fluorescamine positive from the Vydac C-18 column elution shown in Figure 19.

Figure 21 is the elution pattern through a YMC ODS column of a tryptic digest of fractions 36 and 37 of the Mono Q Chromatography.

Figure 22 shows a summary of the sequences obtained from CNBr fragments of CLMF produced in Example 6.

Figure 23 shows the reverse-phase HPLC of peptide fragments of CLMF produced according to Example 6.

Figure 24 is an SDS PAGE of pure CLMF and "free" unassociated 40 kDa subunit of CLMF purified by Affinity Chromatography according to Example 7.

Figure 25 shows the DNA and deduced amino acid sequences for the 40 kDa subunit of CLMF.

Figure 26 shows the cDNA sequence and deduced amino acid sequence for the human 35 kDa CLMF subunit.

Figure 27 shows the inhibition of CLMF bioactivity by serum from CLMF immunized and control rats.

Figure 28 shows SDS PAGE analysis of immunoprecipitation of ^{125}I -CLMF by monoclonal antibodies 4A1 (1), 4D1 (2), 8E3 (3), 9C8 (4) and control (5) and by immune rat serum (6 and 8) and normal rat serum (7 and 9).

5

Figure 29 shows the immunodepletion of CLMF bioactivity (TGF activity) by monoclonal anti-CLMF antibodies (a-CLMF).

Figure 30 shows immunodepletion of CLMF bioactivity (LAK induction activity) by monoclonal anti-CLMF antibodies (a-CLMF).

10

Figure 31 shows Western blot analysis of the reactivity of monoclonal and rat polyclonal anti-CLMF antibodies with CLMF 75 kDa heterodimer.

15

Figure 32 shows Western blot analysis of the reactivity of monoclonal and rat polyclonal anti-CLMF antibodies with CLMF 40 kDa subunit.

20

Figure 33 shows the binding of ^{125}I -CLMF to PHA-activated peripheral blood lymphocyte (PBL) lymphoblasts.

25

Figure 34 shows the inhibition of ^{125}I -CLMF binding to PHA-activated PBL blast cells by rat anti-CLMF serum. The data are expressed as amount (%bound) of ^{125}I -CLMF binding to the cells in the presence of the indicated concentrations of serum when compared to the total specific binding in the absence of serum.

30

Figure 35 shows the inhibition of ^{125}I -CLMF binding to PHA-activated PBL blast cells by monoclonal antibody supernatants. The data are expressed as % inhibition of ^{125}I -CLMF binding to the cells in the presence of a 1:1 dilution of supernatant when compared to the total specific binding in the absence of antibody supernatant.

Figure 36 shows the inhibition of ^{125}I -CLMF binding to PHA-activated PBL blast cells by various concentrations of purified monoclonal antibodies. The data are expressed as the amount (% cpm bound) of ^{125}I -CLMF binding to the cells in the presence of the indicated concentrations of antibody when compared to the total specific binding in the absence of antibody.

Figure 37 shows Western blot analysis of the reactivity of a rabbit polyclonal anti-CLMF antibody with 75 kDa CLMF (nonreduced) and with 35 kDa CLMF subunit (reduced). The antibody was prepared against a synthetic peptide fragment of the 35 kDa CLMF subunit.

Figure 38 shows the effect of IL-12 on IgE and IFN- γ production by IL-4-stimulated PBMC.

A. PBMC were cultured for 12 days in the presence of 10 ng/ml of IL-4 and increasing concentrations of IL-12 (0.1 to 100 pM). Shown are the mean \pm 1 SEM of three experiments; *significantly different from control without IL-2 at $P < 0.01$, Student's t test.

B. Cells were cultured in the presence of IL-4 and IL-12 (60 pM), with or without anti-IL-12 mAbs (a mixture of antibodies 4A1 and 20C2, each at 10 $\mu\text{g/ml}$).

Fig 39 shows that IL-12 suppresses the accumulation of productive but not germ-line C_e transcripts. Total RNA was extracted from PBMC cultured for 10 days with 10 ng/ml of IL-4; in the absence (lanes 1, 3 and 5) or in the presence of 60 pM IL-12 (lanes 2, 4, and 6). Northern blot was performed as described. The membrane was hybridized with ^{32}P -labeled probes specific for the germ-line transcript (lanes 1 and 2) or for the C_{e1} - C_{e2} region binding to both germ-line and mature C_e transcripts (lanes 3 and 4). Equal loading was assessed by methylene blue staining of the ribosomal RNA (lanes 5 and 6).

DETAILED DESCRIPTION OF THE INVENTION

5 All publications mentioned herein, both supra and infra, are hereby incorporated herein by reference.

The CLMF (or IL-12) active proteins of the present invention include the homogenous natural CLMF protein as well as CLMF proteins which contain a biologically active fragment of the amino acid sequence of natural CLMF and CLMF proteins which contain the amino acid sequence of natural CLMF together with other amino acids. The proteins of this invention have the biological activity of CLMF as measured by standard assays such as T-cell growth factor assay as described in Example 9.

15 The protein of the present invention also include proteins which contain analogous amino acid sequences to CLMF or its CLMF active fragments. Such analogues are proteins in which one or more of the amino acids of natural CLMF or its fragments have been replaced or deleted without eliminating CLMF activity. Such analogues may be produced by known methods of peptide chemistry or by known methods of recombinant DNA technology, such as planned mutagenesis. The CLMF biological activity of all of the proteins of the present invention including the fragments and analogues may be determined by using the standard T-cell growth factor assay such as described in Example 9.

In accordance with this present invention, natural CLMF is obtained in pure form. The amino acid sequence of this protein is depicted in Figures 23 and 24. From the sequence of this protein obtained in accordance with this invention, biologically active analogues and fragments of this CLMF protein can be obtained. These biologically active proteins may be produced biologically through standard recombinant technology or may be chemically synthesized using the sequence described above and an amino acid synthesizer or manual synthesis using chemical conditions well

- known to form peptide bonds between selected amino acids. In this manner, these analogues, fragments and proteins which contain the amino acid sequence of CLMF together with other amino acids can be produced. All of these proteins may then be tested for CLMF activity.

Cytotoxic Lymphocyte Maturation Factor of the present invention was isolated and purified as follows:

10 Production of Supernatant Liquid Containing CLMF.

Human NC-37 B lymphoblastoid cells ATCC CCL 214 (American Type Culture Collection, Rockville, MD) were used for production of CLMF. These cells were maintained by serial passage in RPMI 1640 medium supplemented with 5% heat-inactivated (56°C, 30 min.) fetal bovine serum, 2 mM L-glutamine, 100 units/ml penicillin, and 100 µg/ml streptomycin (all obtained from GIBCO Laboratories, Grand Island, NY).

- 20 Higher producer sublines of NC-37 cells were derived by limiting dilution cloning in liquid microcultures. Each well of three Costar 3596® microplates (Costar Co., Cambridge, MA) received 100 µl of a cell suspension containing five NC-37 cells/ml. The medium used for the cloning was a 1:1 mixture of fresh passage medium and filtered, conditioned medium from stock cultures of the parent NC-37 cells. One week and two weeks after culture initiation each of the microcultures was fed with 50 µl of the 1:1 mix of fresh and conditioned medium. Between 3 and 4 weeks after culture initiation the contents of wells containing clones of NC-37 cells were harvested and passed into larger cultures.

- When the number of cells in a given subline exceeded 1.4×10^6 , one million cells were stimulated to produce CLMF in 1 ml cultures containing 3 ng/ml phorbol 12-myristate 13-acetate (PMA) (Sigma Chemical Co., St. Louis, MO) and 100 ng/ml calcium ionophore A23187 (Sigma). Supernatants were harvested from the cultures

after 2 days, dialyzed against about 50 volumes of Dulbecco's phosphate buffered saline (Gibco) using SPECTROPOR® #1 tubing (Fisher Scientific) overnight with one change of buffer and then for 4 hours against 50 volumes of RPMI 1640 medium with 50 µg/ml of gentamicin (both from Gibco) and tested for CLMF by means of the T cell growth factor assay (see below). Three sublines, NC-37.89, NC-37.98, and NC-37.102, were identified which routinely produced CLMF at titers ≥ 4 times the titers produced by the parent NC-37 cell line. Since cells from these three sublines produced CLMF at similar titers (≥ 800 units/ml), culture supernatants derived from the three sublines were pooled for use as starting material for the purification of CLMF.

Bulk production of CLMF was carried out in roller bottle cultures on a roller apparatus set at about 38 rpms (Wheaton Cell Production Roller Apparatus Model II, Wheaton Instruments, Millville, NJ). Cell suspensions were prepared containing $1-1.5 \times 10^6$ NC-37.89, NC-37.98 or NC-37.102 cells/ml of RPMI 1640 supplemented with 1% Nutridoma-SP (Boehringer Mannheim Biochemicals, Indianapolis, IN), 2 mM L-glutamine, 100 units/ml penicillin, 100 µg/ml streptomycin, 10 ng/ml PMA and 20-25 ng/ml calcium ionophore A23187. Two hundred-fifty to three hundred fifty ml aliquots of the cell suspensions were added to Falcon 3027 tissue culture roller bottles (Becton Dickinson, Lincoln Park, NJ) which had been gassed with a mixture of 5% CO₂, 95% air. The roller bottles were then capped tightly and incubated at 37°C with continuous rolling for three days. At the end of this time, the culture supernatants were harvested. EDTA and phenylmethylsulfonyl fluoride (both from Boehringer Mannheim) were added to the culture supernatants at final concentrations of 1 mM and 0.1 mM, respectively, to retard proteolytic degradation. The supernatants were stored at -4°C until concentration.

Lymphokine Activated Killer (LAK) Cell Induction (L.CI)
Assay.

- Culture supernatants and chromatographic fractions were tested for their ability to synergize with rIL-2 to induce the generation of cytolytic LAK cells as follows. Human peripheral blood mononuclear cells (PBMC) were isolated by the following method. Blood from normal volunteer donors was drawn into syringes containing sufficient sterile preservative-free heparin (Sigma) to give a final concentration of approximately 5 units/ml. The blood was diluted 1:1 with Hanks' balanced salt solution (HBSS) without calcium or magnesium (GIBCO). The diluted blood was then layered over 15 ml aliquots of Ficoll/sodium diatrizoate solution (Lymphocyte Separation Medium, Organon Teknika Corp., Durham, NC) in 50 ml Falcon 2098 centrifuge tubes. The tubes were centrifuged for 30 minutes at room temperature at 500 x g. Following centrifugation, the cells floating on the Ficoll/sodium diatrizoate layer were collected and diluted by mixing with ≥ 2 volumes of HBSS without calcium or magnesium. The resulting cell suspension was then layered over 15 ml aliquots of 20% sucrose (Fisher) in RPMI 1640 medium with 1% human AB serum (Irvine Scientific, Santa Ana, CA) in Falcon 2098 centrifuge tubes. The tubes were centrifuged for 10 minutes at room temperature at 500 x g, and the supernatant fluids were discarded. The cell pellets were resuspended in 5 ml of HBSS without calcium or magnesium; repelleted by centrifugation, and finally resuspended in the appropriate culture medium. Accessory cells were removed from the PBMC by treatment with 5 mM L-glutamic acid dimethyl ester (Sigma) using the same conditions described by Thiele et al. J. Immunol. 131:2282-2290, (1983) for accessory cell depletion by L-leucine methyl ester except that the glutamic acid ester was substituted for the leucine ester.

- The accessory cell-depleted PBMC were further fractionated by centrifugation on a discontinuous Percoll density gradient (Pharmacia, Piscataway, NJ) as described by Wong et al., Cell Immunol. 111:39-54, (1988). Mononuclear cells recovered from the

38, 41, 45, and 58% Percoll layers were pooled and used as a source of LAK cell precursors in the assay. The cells recovered from the Percoll gradient were washed and suspended in tissue culture medium (TCM) composed of a 1:1 mixture of RPMI 1640 and Dulbecco's modified Eagle's medium, supplemented with 0.1 mM nonessential amino acids, 60 µg/ml arginine HCl, 10 mM HEPES buffer, 2 mM L-glutamine, 100 units/ml penicillin, 100 µg/ml streptomycin (all available from GIBCO), 5×10^{-5} M 2-mercaptoethanol (Fisher Scientific, Fair Lawn, NJ), 1 mg/ml dextrose (Fisher), and 5% human AB serum (Irvine Scientific, Santa Ana, CA). These cells were incubated in 24-well tissue culture plates (Costar, Cambridge, MA) in 1 ml cultures (7.5×10^5 cells/culture) to which 10^{-4} M hydrocortisone sodium succinate (Sigma) was added to minimize endogenous cytokine production. Some cultures also received human rIL-2 (supplied by Hoffmann-La Roche) at a final concentration of 5 units/ml and/or supernatants to be assayed for CLMF activity. All cultures were incubated for 3-4 days at 37°C in a humidified atmosphere of 5% CO₂, 95% air.

At the end of this incubation, the contents of each culture were harvested, and the cells were pelleted by centrifugation and resuspended in 0.5 ml of fresh TCM. One tenth ml aliquots of these cell suspensions were mixed with 0.1 ml aliquots of ⁵¹Cr-labelled K562 or Raji cells (both cell lines obtained from the ATCC) and tested for their lytic activity in 5 hour ⁵¹Cr release assays. The method for labelling target cells with ⁵¹Cr and performing the cytolytic assays have been described by Gately et al., [JNCI 69:1245-1254 (1982)]. The percent specific ⁵¹Cr release was calculated as $[(g - c)/(100 - c)] \times 100$, where g is the percentage of ⁵¹Cr released from target cells incubated with lymphocytes and c is the percentage of ⁵¹Cr released spontaneously from target cells incubated alone. The total releasable ⁵¹Cr was determined by lysis of the target cells with 2% sodium dodecyl sulfate; see Gately et al., JNCI 69:1245-1254 (1982). All lymphocyte populations were assayed in quadruplicate for lytic activity.

- LAK Cell Induction Microassay. The microassay for measuring synergy between rIL-2 and CLMF-containing solutions in the induction of human LAK cells was similar to the LAK cell induction assay described above but with the following modifications. Human
- 5 peripheral blood mononuclear cells which had been depleted of accessory cells and fractionated by Percoll gradient centrifugation as described above were added to the wells of Costar 3596 microplates (5×10^4 cells/well). Some of the wells also received rIL-2 (5
- 10 units/ml final concentration) and/or purified CLMF or immunodepleted CLMF-containing solutions. All cultures contained 10^{-4} M hydrocortisone sodium succinate (Sigma) and were brought to a total volume of 0.1 ml by addition of TCM with 5% human AB serum. The cultures were incubated for 3 days at 37°C , after which 0.1 ml of ^{51}Cr -labelled K562 cells (5×10^4 cells/ml in TCM with 5%
- 15 human AB serum) were added to each well. The cultures were then incubated overnight at 37°C . Following this, the cultures were centrifuged for 5 minutes at $500 \times g$, and the supernatant solutions were harvested by use of a Skatron supernatant collection system (Skatron, Sterling, VA). The amount of ^{51}Cr released into each
- 20 supernatant solution was measured with a gamma counter (Packard, Downer's Grove, IL), and the % specific ^{51}Cr release was calculated as described above. All samples were assayed in quadruplicate.

Cytolytic T Lymphocyte (CTL) Generation Assay.

- 25 Methods used for generating and measuring the lytic activity of human CTL have been described in detail in Gately et al., J. Immunol. 136: 1274-1282, 1986, and Wong et al., Cell. Immunol. 111: 39-54, 1988. Human peripheral blood mononuclear cells were isolated from the blood of normal volunteer donors, depleted of accessory cells by
- 30 treatment with L-glutamic acid dimethyl ester, and fractionated by Percoll gradient centrifugation as described above. High-density lymphocytes recovered from the interface between the 45% and 58% Percoll layers were used as responder lymphocytes in mixed
- 35 lymphocyte-tumor cultures (MLTC). CTL were generated in MLTC in 24-well tissue culture plates (Costar #3424) by incubation of Percoll

gradient-derived high density lymphocytes (7.5×10^5 /culture) together with 1×10^5 uv-irradiated HT144 melanoma cells (ATCC, Rockville, MD) or with 5×10^4 gamma-irradiated HT144 melanoma cells in TCM with 5% human AB serum (1.2 ml/culture). For uv-irradiation, HT144 cells were suspended at a density of $1-1.5 \times 10^6$ cells/ml in Hanks' balanced salt solution without phenol red (GIBCO) containing 1% human AB serum. One ml aliquots of the cell suspension were added to 35 x 10 mm plastic tissue culture dishes (Falcon #3001), and the cells were then irradiated ($960 \mu\text{W}/\text{cm}^2$ for 5 min) by use of a 254 nm uv light (model UVG-54 MINERALIGHT[®] lamp, Ultra-violet Products, Inc., San Gabriel, CA). For gamma irradiation, HT144 cells were suspended at a density of $1-5 \times 10^6$ cells/ml in TCM with 5% human AB serum and irradiated (10,000 rad) by use of a cesium source irradiator (model 143, J.L. Shepherd and Associates, San Fernando, CA). Uv- or gamma-irradiated HT144 were centrifuged and resuspended in TCM with 5% human AB serum at the desired cell density for addition to the MLTC. In addition to lymphocytes and melanoma cells, some MLTC received human rIL-2 and/or purified human CLMF at the concentrations indicated in the table. Hydrocortisone sodium succinate (Sigma) was added to the MLTC at a final concentration of 10^{-4} M (cultures containing uv-irradiated melanoma cells) or 10^{-5} M (cultures containing gamma-irradiated melanoma cells) to suppress endogenous cytokine production [S. Gillis et al., J. Immunol. 123: 1624-1631, 1979] and to reduce the generation of nonspecific LAK cells in the cultures [L.M. Muul and M.K. Gately, J. Immunol. 132: 1202-1207, 1984]. The cultures were incubated at 37°C in a humidified atmosphere of 5% CO₂ in air for 6 days. At the end of this time, lymphocytes from replicate cultures were pooled, centrifuged, resuspended in 1.2 ml TCM containing 5% human AB serum, and tested for their ability to lyse HT144 melanoma cells; and, as a specificity control, K562 erythroleukemia cells in overnight ⁵¹Cr release assays.

Melanoma cells and K562 cells were labeled with ⁵¹Cr sodium chromate as described by Gately et al. (JNCI 69: 1245-1254, 1982). Likewise, measurement of lymphocyte-mediated lysis of ⁵¹Cr-

labeled melanoma cells was performed in a manner identical to that described by Gately et al. (ibid.) for quantitating lysis of glioma target cells. For assaying the lysis of ^{51}Cr -labeled K562 cells, 0.1 ml aliquots of lymphocyte suspensions were mixed with 25 μl aliquots of ^{51}Cr -labeled K562 (2×10^5 cells/ml in TCM with 5% human AB serum) in the wells of Costar 3696 "half-area" microtest plates. After overnight incubation at 37°C , the plates were centrifuged for 5 min at $1400 \times g$, and 50 μl of culture medium was aspirated from each well. The amount of ^{51}Cr in each sample was measured with a gamma counter (Packard), and the % specific ^{51}Cr release was calculated as described above. All assays were performed in quadruplicate, and values in the table represent the means ± 1 S.E.M. of replicate samples.

15 T cell growth factor (TGF) assay.

The ability of culture supernatants and chromatographic fractions to stimulate the proliferation of PHA-activated human T lymphoblasts was measured as follows. Human PBMC were isolated by centrifugation over discontinuous Ficoll and sucrose gradients as described above for the LCI assay. The PBMC (5×10^5 cells/ml) were cultured at 37°C in TCM containing 0.1% phytohemagglutinin-P (PHA-P) (Difco Laboratories, Detroit, MI). After 3 days, the cultures were split 1:1 with fresh TCM, and human rIL-2 was added to each culture to give a final concentration of 50 units/ml. The cultures were then incubated for an additional 1 to 2 days, at which time the cells were harvested, washed, and resuspended in TCM at 4×10^5 cells/ml. To this cell suspension was added heat-inactivated goat anti-human rIL-2 antiserum (final dilution: 1/200) to block any potential IL-2-induced cell proliferation in the assay. This antiserum, which was provided by R. Chizzonite, Molecular Genetics Department, Hoffmann-La Roche, was shown to cause 50% neutralization of 2 units/ml rIL-2 at a serum dilution of 1/20,000. An equally functional anti-human rIL2 antibody can be obtained from the Genzyme Co., Boston, MA.

Fifty μ l aliquots of the cell suspension containing anti-IL-2 antiserum were mixed with 50 μ l aliquots of serial dilutions of culture supernatants or chromatographic fractions in the wells of Costar 3596 microplates. The cultures were incubated for 1 day at 37°C in a humidified atmosphere of 5% CO₂ in air, and 50 μ l of ³H-thymidine (New England Nuclear, Boston, MA), 10 μ Ci/ml in TCM, were then added to each well. The cultures were further incubated overnight. Subsequently, the culture contents were harvested onto glass fiber filters by means of a cell harvester (Cambridge Technology Inc., Cambridge, MA), and ³H-thymidine incorporation into cellular DNA was measured by liquid scintillation counting. All samples were assayed in triplicate.

In purifying CLMF it was necessary to define units of activity in order to construct chromatographic elution profiles and to calculate the percent recovery of activity and the specific activity of the purified material. To do this, a partially purified preparation of human cytokines produced by coculturing PHA-activated human PBMC with NC-37 cells was used as a standard. Several dilutions of this preparation, which was assigned an arbitrary titer of 2000 units/ml, were included in each TGF or LAK induction assay. The results obtained for the standard preparation were used to construct a dose-response curve from which could be interpolated units/ml of activity in each unknown sample at the dilution tested. Multiplication of this value by the dilution factor yielded the activity of the original sample expressed in units/ml.

For antibody neutralization studies, the TGF assay was modified as follows: Twenty-five μ l aliquots of CLMF-containing medium were mixed with 50 μ l aliquots of serial dilutions of antiserum or antibody solutions in the wells of COSTAR 3596® microplates. The mixtures were incubated for 30 min. at 37°C, and 25 μ l aliquots of a suspension of PHA-activated lymphoblasts (8×10^5 /ml in TCM plus 1:100 anti-rIL-2) were then added to each well. The cultures were

further incubated, pulsed with ^3H -thymidine, harvested, and analyzed for ^3H -thymidine incorporation as described above.

Natural killer (NK) cell activation assay.

5

Purified CLMF was tested for its ability to activate NK cells when added alone or in combination with rIL-2 as

follows: Human PBMC were isolated by centrifugation over discontinuous Ficoll and sucrose gradients as described above and

- 10 were suspended in RPMI 1640 medium supplemented with 10% heat-inactivated fetal bovine serum, 100 units/ml penicillin, 100 $\mu\text{g/ml}$ streptomycin, and 2 mM L-glutamine. The PBMC were incubated overnight at 37°C in 1 ml cultures (5×10^6 cells/culture) together with rIL-2 and/or purified CLMF at various concentrations.
- 15 After 18-20 hours, the contents of the cultures were harvested and centrifuged, and the cells were resuspended in the same medium used for the overnight cultures. The cytolytic activity of the cultured PBMC was then assessed in ^{51}Cr release assays as described above.

- 20 Purification and Characterization of Cytotoxic Lymphocyte Maturation Factor (CLMF)

Concentration of cell supernatant solutions

- 25 Crude human CLMF supernatant solutions prepared from several batches of induced NC-37 cells were pooled and concentrated 30-fold using the Pellicon Cassette System (30,000 NMWL PTTK0005; Millipore Corp., Bedford, MA). After concentrating to the desired volume, a buffer exchange was performed with 10 mM MES, pH adjusted to 6.0 with 10N NaOH. The concentrate was centrifuged
- 30 at 10,000 x g for 10 min at 4°C and the precipitate discarded.

Ion-Exchange Chromatography on NuGel P-SP Column

- The concentrated supernatant solution was applied at a flow rate of 120 ml/hr to a NuGel P-SP (Separation Industries, Metuchen, NJ) column (5 x 5 cm), equilibrated in 10mM MES, pH 6.0. The column was washed until baseline absorbance monitoring at 280 nm was obtained. Absorbed proteins were then eluted with a 500 ml salt gradient from 0 to 0.5M NaCl/10mM MES, pH 6.0 at a flow rate of 2 ml/min. Aliquots of fractions were assayed for TGF activity. Fractions containing TGF activity were pooled and dialyzed (Spectra/Por 7, Fisher Scientific) against 50 vol 20mM Tris/HCl, pH 7.5.

Dye-Affinity Chromatography on Blue B-Agarose Column

- The dialyzed sample was centrifuged at 10,000 x g for 10 min at 4°C and the precipitate discarded. The supernatant solution was applied at a flow rate of 20 ml/hr to a Blue B-Agarose (Amicon, Danvers, MA) column (2.5 x 10 cm) equilibrated in 20mM Tris/HCl, pH 7.5. The column was washed with this same buffer until baseline absorbance monitoring at 280 nm was obtained. Absorbed proteins were then eluted with a 500 ml salt gradient from 0 to 0.5M NaCl/20mM Tris/HCl, pH 7.5 at a flow rate of 15 ml/hr. Aliquots of fractions were assayed for TGF activity. Fractions containing TGF activity were pooled and dialyzed (Spectra/Por 7, Fisher Scientific) against 100 vol 20mM Tris/HCl, pH 7.5.

Ion-Exchange Chromatography on Mono Q Chromatography

- The dialyzed sample was filtered through a 0.45 µm cellulose acetate filter (Nalgene Co., Rochester, NY) and the filtrate applied at a flow rate of 60 ml/hr to a Mono Q HR 5/5 (Pharmacia LKB Biotechnology, Inc., Piscataway, NJ) column (5 x 50mm) equilibrated in 20mM Tris/HCl, pH 7.5. The column was washed with this same buffer until baseline absorbance monitoring at 280 nm was obtained. Absorbed proteins were then eluted with a 1 hr linear salt gradient

from 0 to 0.25M NaCl/20mM Tris/HCl, pH 7.5 at a flow rate of 60 ml/hr. Aliquots of fractions were assayed for TGF activity and protein purity was assessed without reduction by SDS-PAGE [Laemmli, U.K. (1970) *Nature* (London) **227**:680-685] using 12% slab gels. Gels were silver stained [Morrissey, *Anal. Biochem.* **117**:307-310] to visualize protein. Fractions which did not contain a single band were further purified by reversed-phase HPLC.

Reversed-Phase HPLC

- 10 The chromatographic system has been described previously by Stern, A.S. and Lewis, R.V. (1985) in *Research Methods in Neurochemistry*, Eds. Marks, N. and Rodnight, R. (Plenum, New York) Vol. 6, 153-193. An automated fluorescence detection system using fluorescamine (Polysciences, Inc., Warrington, PA) monitored the protein in the column effluents [Stein, S. and Moschera, J. (1981) *Methods Enzymol.* **78**:435-447]. Reversed-phase HPLC was carried out using Vydac C18 or diphenyl columns (4.6 x 20mm, The Sep/a/rations Group, Hesperia, CA). Proteins were eluted with an acetonitrile gradient in 0.1% TFA.

20 Protein Analysis

- Amino acid analysis was performed on an instrument which used post-column reaction with fluorescamine for detection [Pan, Y.-C.E., and Stein, S. (1986) in *Methods of Protein Microcharacterization* (Shively, J.E., Ed.), pp. 105-119, Humana Press, Clifton, NJ].

- Sequence analysis was performed using an Applied Biosystems Inc. Model 470A gas phase sequencer (Foster City, CA) [Hewick, R.M., Henkappeler, M.W., Hood, L.E., and Dreyer, W.J. (1981) *J. Biol. Chem.* **256**:7990-7997]. Phenylthiohydantoin (PTH) amino acid derivatives were identified "on-line" with an ABI Model 120A PTH analyzer.

Cloning of the CLMF Gene

As used herein, the term "CLMF polynucleotide containing a sequence corresponding to a cDNA" means that the CLMF
5 polynucleotide contains a sequence which is homologous to or complementary to a sequence in the designated DNA; the degree of homology or complementary to the cDNA will be approximately 50% or greater, will preferably be at least about 70%, and even more preferably will be at least about 90%. The correspondence between
10 the CLMF sequences and the cDNA can be determined by techniques known in the art, including, for example, a direct comparison of the sequenced material with the cDNAs described, or hybridization and digestion with single strand nucleases, followed by size determination of the digested fragments.

15 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA and immunology, which are within the skill of the art. Such techniques are explained fully in the
20 literature. See e.g., Maniatis, Fritsch & Sambrook, Molecular Cloning; A Laboratory Manual (1982); DNA Cloning, Volumes I AND II (D.N. Glover ed. 1985); Oligonucleotide Synthesis (M.J. Gait ed. 1984); Nucleic Acid Hybridization (B.D. Hames & S.J. Higgins eds. 1984); Transcription and Translation (B.D. Hames & S.J. Higgins eds. 1984);
25 Animal Cell Culture (R.I. Freshney ed. 1986); Immobilized Cells and Enzymes (IRL Press, 1986); B. Perbal, A. Practical Guide to Molecular Cloning (1984); the series, Methods in Enzymology (Academic Press, Inc.); Gene Transfer Vectors for Mammalian Cells (J.H. Miller and M.P. Calos eds. 1987, Cold Spring Harbor Laboratory), Methods in
30 Enzymology Vol. 154 and Vol. 155 (Wu and Grossman, and Wu, eds., respectively), Mayer and Walker, eds. (1987), Immunochemical Methods in cell and Molecular Biology (Academic Press, London), Scopes, (1987), Protein Purification: Principles and Practice, Second Edition (Springer-Verlag, N.Y.), and Handbook of Experimental
35 Immunology, Volumes I-IV (D.M. Weir and C.C. Blackwell eds 1986).

The DNA sequences and DNA molecules of the present invention may be expressed using a wide variety of host/vector combinations. For example, useful vectors may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences, such as various known derivatives of SV-40 and known bacterial plasmids, e.g., plasmids from E. coli including pCR1, pBR322, pMB9 and RP4, phage DNAs, e.g., the numerous derivatives of phageg, and other DNA phages, e.g., M13 and other Filamentous single-stranded DNA phages, vectors useful in yeasts, such as the 2m plasmid, vectors useful in eukaryotic cells, such as vectors useful in animal cells, such as those containing SV-40 adenovirus and retrovirus derived DNA sequences and vectors derived from combinations of plasmids and phage DNA's, such as plasmids which have been modified to employ phage DNA or other derivatives thereof.

Such expression vectors are also characterized by at least one expression control sequence that may be operatively-linked to the CLMF DNA sequence inserted in the vector in order to control and to regulate the expression of that cloned DNA sequence. Examples of useful expression control sequences are the lac system, the trp system, the tac system, the trc system, major operator and promoter regions of phage λ , the control region of fd coat protein, the glycolytic promoters of yeast, e.g., the promoter for 3-phosphoglycerate kinase, the promoters of yeast acid phosphatase, e.g., Pho 5, the promoters of the yeast α -mating factors, and promoters derived from polymoma, adenovirus, retrovirus, and simian virus, e.g., the early and late promoters on SV40, and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells and their viruses or combinations thereof.

Among such useful expression vectors are vectors that enable the expression of the cloned CLMF-related DNA sequences in eukaryotic hosts, such as animal and human cells [e.g., P. J. Southern and P. Berg, J. Mol. Appl. Genet., 1, pp. 327-41 (1982); S. Subramani et al., Mol. Cell. Biol., 1, pp. 854-64 (1981); R. J. Kaufmann and P. A. Sharp, Mol. Cell. Biol., 159, pp. 601-64 (1982) S. I. Scahill et al.,

"Expression and Characterization of The Product Of A Human Immune Interferon DNA Gene in Chinese Hamster Ovary Cells", Proc. Natl. Acad. Sci. U.S.A., 80, pp. 4654-59 (1983); G. Urlaub and L. A. Chasin, Proc. Natl. Acad. Sci. USA, 77, pp. 4216-20 (1989)].

5

Furthermore, within each specific expression vector, various sites may be selected for insertion of the CLMF-related DNA sequences of this invention. These sites are usually designated by the restriction endonuclease which cut them. They are well
10 recognized by those of skill in the art. It is, of course to be understood that an expression vector useful in this invention need not have a restriction endonuclease site for insertion of the chosen DNA fragment. Instead, the vector could be joined to the fragment by alternative means. The expression vector, and in particular the
15 site chosen therein for insertion of a selected DNA fragment and its operative linking therein to an expression control sequence, is determined by a variety of factors, e.g., number of sites susceptible to a particular restriction enzyme, susceptibility of the protein to proteolytic degradation by host cell enzymes, contamination of the
20 protein to be expressed by host cell proteins difficult to remove during purification; expression characteristics, such as the location of start and stop codons relative to the vector sequences, and other factors recognized by those of skill in the art. The choice of a vector and an insertion site for a DNA sequence is determined by a balance
25 of these factors, not all selections being equally effective for a given case.

Not all host/expression vector combinations function with equal efficiency in expressing the DNA sequences of this invention or in
30 producing the CLMF polypeptides of this invention. However, a particular selection of a host/expression vector combination may be made by those of skill in the art after due consideration of the principles set forth herein without departing from the scope of this invention. For example, the selection should be based on a balancing
35 of a number of factors, These include, for example, compatibility of the host and vector, toxicity of the proteins encoded by the DNA

sequence to the host, ease of recovery of the desired protein, expression characteristics of the DNA sequences and the expression control sequences operatively linked to them, biosafety, costs and the folding, form or any other necessary post-expression modifications of the desired protein.

The CLMF produced by fermentation of the prokaryotic and eukaryotic hosts transformed with the DNA sequences of this invention can then be employed in the LAK cell and T cell activator and antitumor compositions and methods of the present invention.

The CLMF of the present invention can also be analyzed to determine their active sites for producing fragments or peptides, including synthetic peptides, having the activity of CLMF. Among the known techniques for determining such active sites are x-ray crystallography, nuclear magnetic resonance, circular dichroism, UV spectroscopy and site specific mutagenesis. Accordingly, these fragments may be employed in methods for stimulating T-cells or LAK cells.

Administration of the polypeptides, or perhaps peptides derived or synthesized from them or using their amino acid sequences, or their salts or pharmaceutically acceptable derivatives thereof, may be via any of the conventionally accepted modes of administration of agents which exhibit antitumor activity. These include parenteral, subcutaneous, intravenous, or intralesional administration.

The preferred form of administration depends on the intended mode of administration and therapeutic application. The compositions also will preferably include conventional pharmaceutically acceptable carriers and may include other medicinal agents, carriers, adjuvants, excipients, etc., e.g., human serum albumin or plasma preparations. Preferably, the compositions of the invention are in the form of a unit dose and will usually be administered one or more times a day.

Methods for Assay of Monoclonal Antibodies

- Purification of CLMF and Labelling of CLMF with ^{125}I CLMF was partially purified from cell supernatants harvested from human peripheral blood lymphocytes (PBLs) or NC-37 cells as described previously. Partially purified CLMF was labelled with ^{125}I by a modification of the Iodogen method (Pierce Chemical Co.). Iodogen (Pierce Chemical Co.) was dissolved in chloroform at a concentration of 0.5 mg/ml and 0.1 ml aliquots were added to 12x75 borosilicate glass tubes. The chloroform was evaporated under a stream of nitrogen and the Iodogen was dried in the center of the bottom of the glass tube. The coated tubes were stored in a dessicator at room temperature (RT) under vacuum. For radiolabeling, 0.5-1.0 mCi ^{125}I -Na (Amersham) was added to an Iodogen coated tube containing 0.50 ml of Tris-Iodination Buffer (25 mM Tris-HCl pH 7.5, 0.4 M NaCl, 1 mM EDTA) and incubated for 4 minutes at RT. The activated ^{125}I solution was transferred to a 1.5 ml tube containing 0.05-0.1 ml CLMF (approximately 5 μg in 0.125 M NaCl, 20 mM Tris-HCl pH 7.5) and the reaction was further incubated for 8 minutes at RT. At the end of the incubation, 0.05 ml of Iodogen Stop Buffer (10 mg/ml tyrosine, 10% glycerol in Dulbecco's phosphate buffered saline (PBS) pH 7.4) was added and reacted for 30 seconds. The mixture was then diluted with 1.0 ml Tris-Iodination Buffer and applied to a BioRad BioGel P10DG (BioRad Laboratories) desalting column for chromatography. The column was eluted with Tris-Iodination Buffer and fractions (1 ml) containing the peak amounts of labelled protein were combined and diluted to 1×10^8 cpm/ml with 25% gelatin in Tris-Iodination buffer. The TCA precipitable radioactivity (10% trichloroacetic acid final concentration) was typically in excess of 95% of the total radioactivity. The radiospecific activity ranged from 6000 cpm/fmol to 10,000 cpm/fmol.

- Immunodepletion of CLMF Hybridoma culture supernatants or purified monoclonal antibodies were tested for their ability to immunodeplete CLMF as follows: Goat anti-rat IgG-agarose beads (Sigma Chemical Co., St. Louis, MO) were washed three times with 10

- ml of PBS (Gibco) supplemented with 1% bovine serum albumin (BSA) (Sigma) (PBS/BSA solution). After washing, the beads were resuspended in PBS/BSA at a final concentration of 50% vol/vol. Aliquots (0.2 ml) of the bead suspension were added to 1.5 ml
- 5 Eppendorf tubes, together with the indicated amounts of monoclonal antibodies or hybridoma supernatant solutions. The volume of each mixture was brought to 1.4 ml by the addition of hybridoma maintenance medium [Iscove's modified Dulbecco's medium (IMDM) with 0.1% fetal bovine serum (FBS), 10% Nutridoma-SP (Boehringer-
- 10 Mannheim), and 2 mM L-glutamine], and the mixtures were then incubated for 2 hours at room temperature on a hematology/chemistry mixer. Following this incubation, the tubes were centrifuged in a Beckman microfuge 12 (1.5 minutes at setting 5), and the supernatants were discarded. The beads were again
- 15 washed three times with PBS/BSA and then resuspended in 1 ml of tissue culture medium (TCM) containing 5% human AB serum and the indicated concentration of purified human CLMF. The tubes were subsequently incubated overnight at 4°C on the mixer. Following this, the beads were removed by centrifugation in the microfuge, and
- 20 the resulting immunodepleted supernatant solutions were assayed for residual CLMF activity in the TGF assay or in the microassay for LAK cell induction.

- Immunoprecipitation Assay. For the immunoprecipitation
- 25 reaction, 0.05 to 0.5 ml of hybridoma supernatant, diluted antisera or purified IgG was added to a 1.5 ml microfuge tube containing 0.1 ml of a 50% suspension of goat-anti-rat IgG coupled to agarose (Sigma Chemical Co.). The assay volume was brought up to 0.5 ml with RIPA Buffer (50 mM NaPO₄ pH 7.5, 150 mM NaCl, 1% Triton-X 100, 1%
- 30 Deoxycholic acid, 0.1% SDS, 1% BSA, and 5 mM EDTA) and the mixture was incubated on a rotating mixer for 2 hours at RT. The beads were pelleted by centrifugation for 1 minute at 12,000 x g and then resuspended in 1 ml RIPA Buffer containing 125I CLMF (1 x 10⁵ cpm). The mixture was then incubated on a rotating mixer for
- 35 16 hours at 4°C. Following this incubation, the beads were pelleted by centrifugation and washed 2X in RIPA without BSA. The beads

- were then washed 1X with .125 M Tris-HCl pH 6.8 and 10% glycerol. The ^{125}I -CLMF bound to the solid phase antibodies was released by adding 10 μl of 2X Laemmli Sample Buffer with and without 5% β -mercaptoethanol and heating for 3 minutes at 95°C. The
- 5 immunoprecipitated ^{125}I -CLMF was analyzed by SDS-PAGE on a 10% or 12% polyacrylamide gel and visualized by autoradiography.

- CLMF Receptor Binding Assay. The ability of hybridoma supernatant solutions, purified IgG or antisera to inhibit the binding
- 10 of ^{125}I -CLMF to PHA-activated human T lymphoblasts was measured as follows: 0.1 ml aliquots of serial dilutions of culture supernatants, purified IgG or antisera were mixed with 0.025 ml aliquots of Binding Buffer (RPMI-1640, 5% FBS, 25 mM HEPES pH 7.4) containing ^{125}I -CLMF (1×10^5 cpm). The mixture was incubated on
- 15 an orbital shaker for 1 hour at RT, then 0.025 ml of activated blasts (5×10^7 cells/ml) was added to each tube. The mixture was further incubated for 1 hour at RT. Non-specific binding was determined by inclusion of 10 nM unlabelled CLMF in the assay. Incubations were carried out in duplicate or triplicate. Cell bound radioactivity was
- 20 separated from free ^{125}I -CLMF by centrifugation of the assay contents through 0.1 ml of an oil mixture (1:2 mixture of Thomas Silicone Fluid 6428-R15: A.H. Thomas, and Silicone Oil AR 200: Gallard-Schlessinger) at 4°C for 90 seconds at 10,000xg. The tip containing the cell pellet was excised and cell bound radioactivity
- 25 was determined in a gamma counter.

- SDS Polyacrylamide Gel Electrophoresis (SDS/PAGE) and Western Blotting. Immunoprecipitated ^{125}I -labelled proteins and partially purified CLMF were treated with Laemmli sample buffer (2% SDS,
- 30 125 mM Tris-HCl, pH 6.8, 10% glycerol, .025% bromphenol blue) with and without 5% β -mercaptoethanol, heated at 95°C for 3 minutes and separated by SDS/PAGE on 7.5% or 12% precast gels (BioRad Laboratories). For the immunoprecipitated ^{125}I -labelled proteins, the gels were stained with 0.2% Coomassie Brilliant Blue in 25%
- 35 isopropyl alcohol and 10% acetic acid, destained in 10% methanol and 10% acetic acid, dried and analyzed by autoradiography. For western

blotting, the proteins separated by SDS/PAGE were transferred to nitrocellulose membrane (0.2 μ) for 16 hours at 100 volts in 10 mM Tris-HCl pH 8.3, 76.8 mM glycine, 20% methanol and 0.01% SDS. The nitrocellulose membrane was blocked for 1 hour at 37°C in 3% gelatin, Tris-HCl pH 7.5, 0.15 M NaCl and then probed with hybridoma supernatant solutions or purified antibody diluted in AB buffer (1% bovine serum albumin, 50 mM sodium phosphate pH 6.5, 0.5 M NaCl, 0.05% Tween 20) for 16 hours at 4°C. After washing with wash buffer (PBS, 0.05% Tween 20), the nitrocellulose strips were incubated for 2 hours at room temperature with goat anti-rat IgG antibody coupled to peroxidase (Boehringer Mannheim Biochemicals) diluted in AB buffer. The nitrocellulose membrane was washed with wash buffer and the bound antibody visualized by incubation for 30 minutes at RT with 4-chloro-1-naphthol (0.5 mg/ml in 0.15% H₂O₂, 0.5 M NaCl, 50 mM Tris-HCl, pH 7.5). The reaction was stopped by extensive washing with distilled water.

In order that our invention herein described may be more fully understood, the following examples are set forth: It should be understood that these examples are for illustrative purposes only and should not be construed as limiting this invention in any way to the specific embodiments recited therein.

Example 1

Purification of CLMF

5 CLMF was produced by a subclone of NC-37 lymphoblastoid cells after costimulation with PMA and calcium ionophore A23187. Stored, frozen supernatant solutions from these cells totaling 60 liters were thawed, pooled and concentrated to approximately 1.9
10 liters using the Pellicon Cassette System. To clarify this concentrate, the preparation was centrifuged and the precipitate discarded.

The supernatant solution was applied to a Nu-Gel P-SP column and protein was eluted with a salt gradient (Fig. 1). Peak T cell
15 growth factor (TGF) activity was determined and the active fractions were pooled and dialyzed in order to reduce the salt concentration of the preparation by 50-fold. This material, after centrifugation to remove particulates, was applied to a Blue-B-Agarose column. Protein was eluted with a salt gradient (Fig. 2). Peak TGF activity
20 was determined and the active fractions were pooled and dialyzed in order to reduce the salt concentration of the preparation by 100-fold. This material, after filtration, was applied to a Mono Q column. Protein was eluted with a salt gradient (Fig. 3). Aliquots of fractions were assayed for TGF activity and protein purity of individual
25 fractions was assessed by SDS-PAGE under non-reducing conditions using a 12% slab gel. The gel was silver stained to visualize protein (Fig. 4). Fractions 36 and 37 were of greater than 95% purity and revealed a major band at 75,000 molecular weight. Fractions 38
30 through 41 containing TGF activity, revealed the 75 kDa protein by SDS-PAGE with major contaminants at 55,000 and 40,000 molecular weight.

Therefore, to eliminate these contaminating proteins, fraction 38 of the previous Mono Q chromatography was diluted 1:1 vol/vol with
35 8M urea and pumped onto a Vydac diphenyl column using an enrichment technique. The column was then washed with 5 ml of 0.1% trifluoroacetic acid. Elution of the proteins was accomplished

with a gradient of 0-70% acetonitrile over 7 hrs in 0.1% trifluoroacetic acid (Fig. 5). Aliquots of fractions were assayed for TGF activity. Protein purity of the fractions containing TGF activity was assessed by SDS-PAGE under non-reducing conditions using a 10% slab gel. The gel was silver stained to visualize protein (Fig. 6). Fractions 86 through 90 were of greater than 95% purity and revealed protein of 75,000 molecular weight. Fractions 87 and 88 were pooled and aliquots were analyzed by SDS-PAGE under reducing (in the presence of β -mercaptoethanol) and non-reducing conditions (in the absence of β -mercaptoethanol). Under the reducing conditions, the 75,000 molecular weight CLMF was separated into two subunits of 40,000 and 35,000 daltons (Fig. 7). Thus we may conclude that CLMF is a 75 kDa heterodimer composed of disulfide-bonded 40 kDa and 35 kDa subunits.

The overall purification of CLMF that was achieved is shown in Table 1. The protein content of the Mono Q- and Vydac diphenyl-purified material was calculated on the basis of amino acid analysis. A specific activity of 8.5×10^7 units/mg and 5.2×10^7 units/mg for Mono Q- and Vydac diphenyl-purified material respectively, was obtained. The fact that the diphenyl-purified protein has a slightly lower specific activity than the Mono Q-purified material may be due to inactivation or denaturation of some of the molecules of CLMF in the HPLC elution solvents (i.e., acetonitrile in 0.1% trifluoroacetic acid).

Chemical Characterization

The ability to prepare homogeneous CLMF allowed for the determination of the amino acid composition and partial sequence analysis of the naturally occurring CLMF protein. Between 10 and 20 picomoles of Mono-Q-purified CLMF was subjected to hydrolysis, and its amino acid composition was determined (Table 2). Proline, cysteine and tryptophan were not determined (ND). Quantitation of histidine was not possible due to a large artifact peak, associated with Tris, coeluting with His (*).

Between 5 and 30 picomoles of diphenyl-purified CLMF was subjected to hydrolysis with and without pre-treatment with performic acid. Complete amino acid composition was thus obtained
5 (Table 3) with the exception of tryptophan.

TABLE 1

Step	Volume (ml)	Pooled Activity (U/ml)	Total Units (U)	Pooled Protein (mg/ml)	Total Protein (mg)	Specific Activity (U/mg)
Pooled Cell Supernatants	60,000	2.58×10^3	1.6×10^8	ND	ND	ND
Ultrafiltered Concentrate	1,940	1.57×10^5	3.0×10^8	1.83	3550	8.5×10^4
NUGel P-SP	90	2.00×10^6	1.8×10^8	0.70	63	2.8×10^6
Blue-B-Agarose	45	3.11×10^6	1.4×10^8	0.24	11	1.3×10^7
Mono Q Fraction 37	1	6.40×10^6	6.4×10^6	0.075	0.075	8.5×10^7
Mono Q Fraction 38-42	5	6.90×10^6	3.45×10^7	0.081	0.405	8.5×10^7
Diphenyl Fraction 87+88	1.1	5.74×10^5	5.22×10^5	0.008	0.010	5.2×10^7

00101839.002299

459151

TABLE 2

	<u>Amino Acid</u>	<u>mol %</u>
5	Aspartic acid or asparagine	11.8
	Threonine	7.8
	Serine	8.4
	Glutamic acid or glutamine	14.9
10	Proline	ND
	Glycine	6.2
	Alanine	7.6
	Cysteine	ND
15	Valine	6.9
	Methionine	2.0
	Isoleucine	4.6
	Leucine	9.0
20	Tyrosine	3.7
	Phenylalanine	4.0
	Histidine	*
	Lysine	9.3
	Arginine	5.4
	Tryptophan	ND

TABLE 3

	<u>Amino acid</u>	<u>mol %</u>
5	Aspartic acid or asparagine	10.8
	Threonine	7.2
	Serine	8.9
	Glutamic acid or glutamine	13.1
10	Proline	3.8
	Glycine	4.7
	Alanine	5.9
	Cysteine	2.9
	Valine	6.2
15	Methionine	1.9
	Isoleucine	4.2
	Leucine	9.4
	Tyrosine	3.6
	Phenylalanine	3.7
20	Histidine	1.8
	Lysine	7.7
	Arginine	4.4
	Tryptophan	ND
25		

Amino-terminal sequence determination was attempted by automated Edman degradation on 100 pmol of the Mono-Q-purified CLMF. Data from the first 22 cycles indicated two sequences present, as would be expected from the heterodimeric structure of CLMF. These results may be summarized as follows:

	Cycle	1	2	3	4	5	6	7	8
5	Amino Acid	I/?	W/?	E/L	L/P	K/V	K/A	D/T	V/P
	Cycle	9	10	11	12	13	14	15	16
10	Amino Acid	Y/D	V/P	V/G	E/M	L/F	D/P	W/?	Y/L
	Cycle	17	18	19	20	21	22		
15	Amino Acid	P/H	D/H	A/S	P/Q	G/?	E/?		
20									

Example 2

Determination of the amino-terminal sequence of the 40,000 dalton subunit of CLMF

Stored supernatant solutions from NC-37 cells totaling 39.1 liters were pooled and concentrated to approximately 2.4 liters using the Pellicon Cassette System and stored at -20°C. To clarify this concentrate after thawing, the preparation was centrifuged and the precipitate discarded.

The supernatant solution was applied to a Nu-Gel P-SP column and protein was eluted with a salt gradient (Fig. 8). Peak TGF activity was determined and the active fractions were pooled and dialyzed in order to reduce the salt concentration of the preparation by 50-fold. This material, after centrifugation to remove particulates, was applied to a Blue-B-Agarose column. Protein was eluted with a salt gradient (Fig. 9). Peak TGF activity was

determined and the active fractions were pooled and dialyzed in order to reduce the salt concentration of the preparation by 100-fold. This material, after filtration, was applied to a Mono Q column. Protein was eluted with a salt gradient (Fig. 10). Aliquots of
5 fractions were assayed for TGF activity.

Fractions 39 and 40 of the previous Mono Q chromatography were pooled and diluted 1:1 vol/vol with 8M urea and pumped onto a Vydac diphenyl column using an enrichment technique. The
10 column was then washed with 5 ml of 0.1% trifluoroacetic acid. Elution of the proteins was accomplished with a gradient of 0-70% acetonitrile over 7 hrs in 0.1% trifluoroacetic acid (Fig. 11). Aliquots of fractions were assayed for TGF activity. Protein purity of the fractions containing TGF activity was assessed by SDS-PAGE under
15 reducing (in the presence of β -mercaptoethanol) conditions (Fig. 12). Fractions 94 through 97 contained the 40,000 dalton subunit >90% pure.

Chemical Characterization

The ability to prepare a highly enriched preparation of the 40,000 dalton subunit of CLMF allowed for its partial sequence analysis.

Amino terminal sequence determination was attempted by automated Edman degradation on 20 pmol of the diphenyl- purified 40,000 dalton subunit. The results may be summarized as follows:

10	Cycle	1	2	3	4	5	6	7
	Amino Acid	I	W	E	L	K	K	D
15	Cycle	8	9	10	11	12	13	14
	Amino Acid	V	Y	V	V	E	L	D
20	Cycle	15	16	17	18	19	20	21
	Amino Acid	W	Y	P	D	A	P	G
25	Cycle	22	23					
	Amino Acid	E	M					
30								

With regard to the sequence analysis of 75,000 dalton CLMF (example 1) and the sequence analysis of the 40,000 dalton subunit of CLMF (example 2), one can deduce the amino terminal sequence of the 35,000 dalton subunit of CLMF. The amino terminal sequences of the 35,000 dalton subunit and the 40,000 dalton subunit can be summarized as follows:

10 35,000 dalton subunit:

NH₂ -?-?-Leu-Pro-Val-Ala-Thr(?)⁵-Pro-Asp-Pro-Gly¹⁰

40,000 dalton subunit:

NH₂ ¹ -Ile-Trp-Glu-Leu-⁵ ~~Ly⁵-Ly⁶-Asp~~-Val-Tyr-Val-Val-Glu ¹⁰

25

where ? represents an undetermined or "best-guessed" residue.

30 Determination of Internal Amino Acid Sequence Segments of the 40,000 dalton Subunit of CLMF

CLMF was purified as previously described in Example 1. The 40,000 dalton subunit was separated and purified from the 35,000 dalton subunit by the method described by Matsudaia [Journal of Biological Chemistry **262**, 10035-10038 (1987)]. Fifty micrograms of CLMF (in 500 μ l of 20 mM Tris, pH 7.5; 0.15 M NaCl) was diluted with 200 μ l 2x concentrate Laemmli sample buffer [Nature **227**, 680-685

(1970)]. The sample was concentrated to 400 μ l and disulfide bonds broken by the addition of 18 μ l β -mercaptoethanol followed by exposure to 105°C for 6 minutes.

5 The sample was loaded onto a minigel (1.0 mm thick) containing 12% polyacrylamide and electrophoresed according to Laemmli. After electrophoresis, the gels were soaked in transfer buffer (10 mM 3-[cyclohexylamino]-1-propanesulfonic acid, 10% methanol, pH 11.0) for 5 min to reduce the amount of Tris and
10 glycine. During this time, a polyvinylidene difluoride (PVDF) membrane (Immobilon; Millipore; Bedford, MA) was rinsed with 100% methanol and stored in transfer buffer. The gel, backed with two sheets of PVDF membrane and several sheets of blotting paper, was assembled into a blotting apparatus and electroeluted for 30 min
15 at 0.5 Amps in transfer buffer. The PVDF membrane was washed in deionized H₂O for 5 min. The edge of the blot was excised from the PVDF membrane and stained with 0.1% Coomassie Blue R-250 in 50% methanol for 5 min, and then destained in 50% methanol, 10% acetic acid for 5-10 min at room temperature. The 40,000 dalton stained
20 band was then matched to the corresponding region of the unstained blot and the 40,000 subunit was cut from the unstained PVDF.

 The Coomassie Blue-stained 40,000 dalton subunit was N-terminal sequenced to confirm that the N-terminus matched that
25 previously obtained (see Example 2). By this method, the 40,000 dalton protein was identified as the 40,000 subunit of CLMF.

 Five percent of the PVDF bound 40,000 dalton subunit was analyzed for its amino acid composition (Table 4). The remaining
30 95% of the blotted 40,000 dalton subunit was fragmented with trypsin according to the procedure of Bauw, et al. [Proc. Natl. Acad. Sci. USA 86, 7701-7705 (1989)]. The membrane carrying the protein was cut into pieces of approximately 3 by 3 mm, and collected in an Eppendorf tube. They were then immersed in 300 μ l of a 2%
35 polyvinylpyrrolidone (40,000 dalton) solution in methanol. After 30 min., the quenching mixture was diluted with an equal volume of

distilled water and further incubated for 5-10 min. The supernatant solution was then discarded and the membrane pieces were washed four times with 300 μ l water and once with 300 μ l 100 mM Tris HCl (pH 8.5). Two hundred microliters of this buffer containing 2 μ g of trypsin was added. The sample was shaken and incubated for 4 hr at 37°C. The supernatant solution was then transferred into a second Eppendorf tube and the membrane pieces were further washed once with 100 μ l of 88% (vol/vol) formic acid and three times with 100 μ l of deionized water. All washing solutions were added to the digestion mixture in the second Eppendorf tube. The resultant peptides contained in the pooled digest were separated by narrow bore HPLC (HP1090A, Hewlett Packard) on a YMC C-18 column (2.6 x 50 mm; Morris Plains, NJ).

The above described procedure is shown, in cartoon form, in Figures 13 & 14.

The tryptic peptide map of the digested 40,000 dalton subunit is shown in Figure 15. Peptides were eluted with a linear gradient of acetonitrile. The peaks which were sequenced are numbered according to their fraction number. The amino acid sequence of these peptides is shown in Table 5.

Many tryptic peptides were recovered from all regions of the intact 40,000 dalton subunit (Table 5). The N-terminal hexapeptide (fraction no. 60) was recovered in high yield. The carboxy-terminal peptide (fraction no. 72) was recovered and is the full length of the predicted C-terminal peptide although the last two amino acids were not positively confirmed by sequencing. This is probably due to the fact that Cys and Ser residues are not detected well, especially when they occur at the end of a peptide. Four potential Asn-linked carbohydrate sites are predicted from the cDNA sequence. Two peptides containing two of these sites were sequenced. When peptide 196-208 (fraction no. 70) was sequenced, no peak was detected at residue 200 indicating that this Asn (predicted by the cDNA) is indeed glycosylated. Peptide 103-108

(fraction no. 52) yielded Asn at residue 103. Therefore, this site is not glycosylated.

- 5 An unknown peak seen in the PTH (sequence) analysis of fraction no. 55 was detected at the position corresponding to residue no. 148. The site is predicted to be a Cys residue which is normally not detected by sequence analysis unless it is modified.

- 10 The above PVDF transfer procedure was repeated on a second 50 mg aliquot of CLMF (see Figure 13 & 14 for procedure outline). However, the blotted 40,000 dalton subunit was fragmented with the proteolytic enzyme, *Staphylococcus aureus* V8 protease (Endoproteinase Glu-C, Boehringer Mannheim, Indianapolis, IN). Membrane pieces were digested for 6 hours at 37°C with 20 mg of
- 15 V8. The peptides were extracted with 88% (vol/vol) formic acid and separated on a Phase Separations column (2 x 150 mm, C8 S3, Queensferry, England, UK) (Figure 16). Peptides were eluted with a linear gradient of acetonitrile. The peaks which were sequenced are numbered according to their fraction number. The amino acid
- 20 sequence of these peptides is shown in Table 6.

- Three major peaks of peptide (fraction nos. 47, 54 and 57) containing four peptides were sequenced. All four peptides were from the amino-terminal region of the 40 kDa subunit indicating that
- 25 the N-terminus of the protein is most susceptible to V8-digestion.

Figure 17 summarizes the protein structural determination of the 40,000 dalton subunit of CLMF.

TABLE 4

	<u>Amino Acid</u>	<u>Residue No.</u>	
5	Aspartic acid or asparagine	27.9	(28)
	Threonine	20.7	(23)
	Serine	24.6	(34)
	Glutamic acid or glutamine	44.6	(35)
	Proline	ND	(14)
10	Glycine	16.3	(15)
	Alanine	16.2	(14)
	Cysteine	ND	(10)
	Valine	20.9	(23)
	Methionine	2.5	(2)
15	Isoleucine	10.3	(12)
	Leucine	22.9	(22)
	Tyrosine	12.9	(12)
	Phenylalanine	9.9	(9)
	Histidine	5.2	(5)
20	Lysine	24.5	(26)
	Arginine	12.5	(12)
	Tryptophan	ND	(10)

- 25 Note: The results represent the mean of two analyses. Proline, cysteine, and tryptophan were not determined (ND). Values in parentheses represent the theoretical amino acid composition of the 40,000 dalton subunit based upon the primary structure of the protein deduced from sequence analysis of cloned 40,000 dalton subunit.
- 30

TABLE 5

Tryptic 40kDa CLMF peptides off PVDF

	<u>fraction</u> <u>no.</u>	<u>residue</u> <u>no.</u>	<u>N-terminal sequence</u>
5	52	103-108	N-K-T-F-L-R
	55	139-157	G-S-S-D-P-Q-G-V-T*-G-A-A-T-L-S-A-E-R
10	55 & 57	267-279(?)	V-F-T-D-K-T-S-A-T-V-I-?-R
	57	52-58	T-L-T-I-Q-V-K
15	57	218-228	N-L-Q-L-K-P-L-K-N-S-R
	60	1-6	I-W-E-L-K-K
	67	288-?	A-Q-D-R-Y-Y-S-S
20	67	85-102(?)	K-E-D-G-I-W-S-T-D-I-L-K-D-Q-K-E-P
	70	196-208	L-K-Y-E-?-Y-T-S-S-F-I-(R?)
25	71	85-96(?)	K-E-D-G-I-?-S-T-D-I-L-K
	72	288-306(?)	A-Q-D-R-Y-Y-S-S-S-W-E-?-A-S-V-P-?-?
	78	71-85	(G?)-G-E-V-L-S-H-S-L-L-L-(L?)-H-K-K

TABLE 6

V8 (Glu-C) 40kDa peptides off PVDF

5	<u>fraction no.</u>	<u>residue no.</u>	<u>N-terminal sequence</u>
	47	1-3	I-W-E
10	54	4-12	L-K-K-D-V-Y-V-V-E
	57	13-22	L-D-W-Y-P-D-A-P-G-E
	57	45-59	V-L-G-S-G-K-T-L-T-I-Q-V-K-(E?)
15			

EXAMPLE 4

20 Direct determination of the amino-terminal sequence of the 35,000 dalton subunit of CLMF

SDS-PAGE analysis of the Mono Q fraction 39 from Example 1 under reducing (in the presence of β -mercaptoethanol) and non-reducing (in the absence of β -mercaptoethanol) conditions (Fig. 18) demonstrated that the 40,000 dalton molecular weight "contaminant" is "free" 40,000 dalton CLMF subunit (i.e. unassociated with the 35,000 dalton subunit). The evidence which points to this deduction is that without reduction (lane 1, Fig. 18) mainly 75,000 dalton CLMF is present with some 40,000 dalton protein. After reduction (lane 2, Fig. 18), the 75,000 dalton CLMF is gone yielding the 35,000 dalton subunit and an enriched 40,000 dalton band:

35 Fraction 39 of the previous Mono Q chromatography was reduced in 5% β -mercaptoethanol in the presence of 4 M urea and heated for 5 mins at 95°C. The sample was pumped onto a

Vydac C-18 column using an enrichment technique and the column was then washed with 5 ml of 0.1% trifluoroacetic acid. Elution of the proteins was accomplished with a gradient of 0-70% acetonitrile over 5 hrs in 0.1% trifluoroacetic acid (Fig. 19). Protein purity of the fractions which were fluorescamine positive was assessed by SDS-PAGE under non-reducing conditions using a 10% slab gel. The gel was silver stained to visualize protein (Fig. 20). Fractions 112 through 117 revealed a diffuse band at 35,000 molecular weight which was greater than 95% pure. The 40,000 dalton subunit and any other proteins present in fraction 39 remained bound to the C-18 column. These proteins (including the 40,000 dalton subunit) were finally eluted with a solution of 42% formic acid/40% 1-propanol.

15 Chemical Characterization

The ability to prepare homogeneous 35,000 subunit allowed for the determination of the amino acid composition and partial sequence analysis of the lower molecular weight subunit of the CLMF protein. Approximately 1 μ g of 35 kDa subunit was subjected to hydrolysis, and its amino acid composition was determined (Table 7). Proline, cysteine and tryptophan were not determined (ND).

Amino-terminal sequence determination was attempted by automated Edman degradation on 100 pmol of the C-18 purified 35kD subunit. Data from the first 20 cycles confirmed the sequence obtained by deduction as described in Example 2. Furthermore, the second amino acid was obtained in addition to amino acids 21 thru 26. These results may be summarized as follows:

	Cycle	1	2	3	4	5	6	7
5	Amino Acid	?	N	L	P	V	A	T
	Cycle	8	9	10	11	12	13	14
10	Amino Acid	P	D	P	G	M	F	P
	Cycle	15	16	17	18	19	20	21
15	Amino Acid	?	L	H	H	S	Q	N
	Cycle	22	23	24	25	26		
20	Amino Acid	L	L	R	A	V		
25								

Therefore, the amino terminal sequence of the 35,000 dalton subunit can be summarized as follows:

30 NH₂-7-Asn-Leu-Pro-Val-Ala-Thr-Pro-Asp-Pro-Gly-Met-
15 20 25 26
35 Phe-Pro-?-Leu-His-His-Ser-Gln-Asn-Leu-Leu-Arg-Ala-Val
where ? represents an undetermined residue.

TABLE 7

	<u>Amino Acid</u>	<u>Mol %</u>
5	Aspartic acid or asparagine	10.9
	Threonine	6.7
	Serine	8.3
	Glutamic acid or glutamine	14.9
10	Proline	ND
	Glycine	6.1
	Alanine	7.7
	Cysteine	ND ₂
	Valine	6.3
15	Methionine	2.9
	Isoleucine	4.5
	Leucine	10.9
	Tyrosine	3.2
	Phenylalanine	4.4
20	Histidine	2.3
	Lysine	5.6
	Arginine	5.5
	Tryptophan	ND

25

Example 5Determination of the sequence of a tryptic fragment of CLMF

- 30 **Mono Q** fractions 36 and 37 from Example 1 were pooled (approximately 100 pmol/1.7 ml) and the volume (less 30 μ l-see Example 6) reduced to 200 μ l under a stream of helium. One hundred microliters of 0.1 M ammonium bicarbonate was added. Trypsin (Worthington Biochemical Corp., Freehold, NJ) cleavage was
- 35 performed at a substrate-to-enzyme ratio of 2:1 (w/w) at 37°C for 20 hr. The resultant peptide fragments were reduced and

carboxymethylated. This was accomplished by addition of 160 μ l of 0.1 M Tris-HCl, pH 8.5/6 M guanidine-HCl. The volume was reduced to 200 μ l under a stream of helium, and 4 μ l of dithiothreitol (50 mg/ml) was added. The mixture was incubated at 37°C for 4 hrs.

- 5 After reductive cleavage of the disulfide bonds, [14 C]iodoacetic acid (4 mmol) was added and the resulting solution was incubated in the dark at room temperature for 10 min.

- The resultant peptide fragments were isolated by reversed-phase HPLC (Fig. 21) on an S-5 120 Angstrom ODS column (2.6 x 50 mm, YMC, Inc., Morris Plains, NJ). Peptides were eluted with a 1-propanol gradient in 0.9 M acetic acid, pH adjusted to 4.0 with pyridine. The amino acid sequence of the peptide found in fraction 46 was found to be (by automated Edman degradation): Asp-Ile-Ile-
15 Lys-Pro-Asp-Pro-Pro-Lys.

Example 6

Determination of Internal Amino Acid Sequence Segments of CLMF

- 20 CLMF was purified as previously described in Example 1. Approximately 80 μ g of protein was precipitated with 10% trichloroacetic acid. The precipitate was dissolved in 70% (v/v) aqueous formic acid at room temperature. An approximately 50-fold
25 molar excess over methionine residues of cyanogen bromide (CNBr) in a small volume of 70% formic acid was added, with stirring, and the mixture was incubated in the dark under oxygen-free helium at room temperature for 48 hr. The mixture was diluted with 15 vol of water, divided into two equal portions and dried under a stream of
30 helium. For complete removal of the acid and by-products, the drying was repeated after further addition of water.

- One of the portions (@40 μ g) of fragmented CLMF was dissolved with 50 μ l Laemmli sample buffer [Nature 227, 680-685
35 (1970)] containing 4% β -mercaptoethanol followed by exposure to 105°C for 6 minutes. The sample was loaded into 3 wells of a

minigel (1.0 mm thick) containing 17.5% polyacrylamide and electrophoresed according to Laemmli.

After electrophoresis, the gels were soaked in transfer buffer (10 mM 3-[cyclohexylamino]-1-propanesulfonic acid, 10% methanol, pH 11.0) for 30 min. During this time, a polyvinylidene difluoride (PVDF) membrane (Immobilon; Millipore; Bedford, MA) was rinsed with 100% methanol and stored in transfer buffer. The gel, backed with two sheets of PVDF membrane and sandwiched with blotting paper, was assembled into a blotting apparatus and electroeluted for 30 min. at 0.5 Amps in transfer buffer. The PVDF membrane was washed in deionized H₂O for 5 min and stained with 0.1% Coomassie Blue R-250 in 50% methanol for 5 min, and then destained in 50% methanol, 10% acetic acid for 5-10 min at room temperature. A number of smeared bands were observed.

Five regions of the membrane were excised across the three last lanes containing the CLMF CNBr digest. These regions were sequenced. A summary of the sequences obtained from the CNBr fragments of CLMF is shown on Fig. 22.

The second portion (@40 µg) of fragment CLMF was dissolved in 400-500 µl 88% formic acid containing 6 M guanidine-HCl, 0.1 M Tris/HCl, 0.5 M NaOH, pH 8.0. The sample was pH adjusted to pH 4.0 with formic acid. The peptide fragments were isolated by reversed-phase HPLC (Fig. 23) on a Vydac C₄ column (4.6 x 20 mm, The Sep/ara/tions Group, Hesperia, CA). Peptides were eluted with a 4.5 hr linear gradient of acetonitrile in 0.1% TFA. One of these peaks was sequenced and the amino acid sequence of this peptide was:

Fraction No.	N-Terminal Sequence
47	V-D-A-V-H-K-L-K-Y-E-2-Y-T-S-(S?) -F-F-I-R-D-I-I-K-P-
35	(Starts at residue #190 of 40 kDa subunit)

It is assumed or known that the above sequence is preceded by a Met residue"?" represents a "best-guessed" residue.

Example 7

5

Purification of CLMF and 40,000 dalton subunit using affinity chromatography

10 An affinity chromatography resin was prepared by covalently attaching 7B2 monoclonal antibody to activated agarose. Similarly, the below outlined purification could also be carried out by covalently coupling the antibody to silica or thin microporous membranes. The activated agarose was prepared as follows:

- 15 1. 100 ml Sepharose CL-6B was washed three times with 100 ml H₂O.
2. 100 ml of 1% sodium meta-periodate in H₂O was added to the resin and the suspension shaken at room temperature for 60 min.
- 20 3. The resin was washed with cold H₂O thoroughly.

The covalent attachment of 7B2 to the activated agarose was carried out as follows:

- 25 1. .9 ml of the activated agarose (described above) was suspended in 7 ml of 7B2 (@ 3.9 mg/ml) in phosphate buffered saline, pH 7.4.
2. Added 50.2 mg of cyanoborohydride was added to the gel suspension which was shaken overnight at 4°C.
- 30 3. The gel suspension was filtered and added to 7 ml of 1.0 M ethanolamine, pH 7.0 containing 50.2 mg of cyanoborohydride.

One millimeter of the above described resin (@ 2.6 mg IgG/ml
35 gel) was packed in a column and washed extensively with phosphate buffered saline. Fractions from the Mono Q chromatography

containing the 75 kDa, CLMF protein but with additional major contaminating proteins, were pooled (approx. 3.5×10^6 U TGF activity) and dialyzed extensively against PBS. This preparation was applied to the 7B2-Sepharose column at a rate of 5 ml/hr at room temperature. The column was washed with phosphate buffered saline (pH 7.4) until baseline absorbance monitoring at 280 nm was obtained. Adsorbed proteins were then eluted with 0.2 N acetic acid, 0.15 M NaCl, pH @3. Aliquots of fractions were assayed for TGF activity. Approximately 76% of the starting activity was recovered in the acid eluate.

Protein purity was assessed without reduction by SDS-PAGE [Laemmli, U.K. (1970) Nature (London) 227: 680,685)] using a 10% slab gel. Gels were silver stained [Morrissey, Anal. Biochem. 117:307-310] to visualize protein. The acid eluant contained pure CLMF and the "free" unassociated 40 kDa subunit of CLMF (Fig. 24).

Example 8

Determination of the pI of CLMF

Thirty microliters of the pooled Mono Q fractions 36 and 37 (see Example 5) were spotted onto a precast ampholine PAGplate gel, pH 3.5-9.5 (Pharmacia LKB Biotechnology) to determine the pI of CLMF. Based on pI standard markers, a major band was observed at pI 4.8 and a minor band at pI 5.2. Based on pH determination, the pI of these bands are 4.2 and 4.6 respectively.

Example 9

Biologic activities of purified CLMF

Purified CLMF stimulated the proliferation of human PHA-activated lymphoblasts in the T cell growth factor assay (Table 8). The T cell growth factor activity of the purified CLMF recovered from the Mono

Q column was compared to that of a standard preparation of human lymphokines in five separate experiments, and the specific activity of the purified CLMF was found to be $8.5 \pm 0.9 \times 10^7$ units/mg protein. In one experiment in which purified CLMF obtained from diphenyl HPLC was compared to the standard lymphokine preparation in the TGF assay, a specific activity of 5.2×10^7 units/mg protein was observed. When suboptimal concentrations of purified CLMF and human rIL-2 were tested in combination in the TGF assay, additive proliferation was observed (Table 8), up to the maximum proliferation caused by rIL-2 alone. However, proliferation caused by rIL-2 could be distinguished from proliferation due to CLMF in that the former was totally inhibited in the presence of a neutralizing goat anti-human IL-2 antiserum but the latter was not affected.

The ability of purified CLMF to activate cytotoxic effector cells was examined both in a 4-day LAK cell induction assay and in an overnight NK cell activation assay. In the LCI assay, purified CLMF at concentrations as high as 800 units/ml had little activity in the absence of IL-2 (Table 9). However, CLMF synergized with low concentrations of human rIL-2 in causing LAK cell induction in as much as the lytic activity generated in the presence of both cytokines was significantly greater than the sum of the lytic activities observed in cultures containing either cytokine alone (Table 9). In the presence of rIL-2, purified CLMF was active at concentrations as low as 3 units/ml.

In contrast to the results in the 4-day LAK induction assay, purified CLMF was effective by itself in activating human NK cells in an overnight assay (Table 10). In this assay, CLMF was active at concentrations as low as 1.6 units/ml. When CLMF was tested in combination with human rIL-2, the two cytokines together had, at best, additive effects in enhancing NK activity (Table 10).

In addition to its ability to enhance the lytic activity of nonspecific NK/LAK cells, CLMF also facilitated specific human cytolytic T lymphocyte (CTL) responses in vitro. CLMF increased the specific allogeneic CTL response to weakly immunogenic, gamma-irradiated HT144 melanoma cells (Table 11). In combination with a low concentration of rIL-2, CLMF also facilitated specific allogeneic human CTL responses to uv-irradiated HT144 melanoma cells, which did not elicit any detectable CTL response in the absence of added cytokines (Table 11). The specificity of the cytolytic effector cells generated in these studies was demonstrated by their ability to cause substantial lysis of ⁵¹Cr-labeled HT 144 melanoma cells but little or no lysis of K562 cells. In contrast, LAK cells which were generated in the same experiments by incubating low density lymphocytes with rIL-2 in the absence of hydrocortisone lysed the K562 cells to a much greater extent than HT144 melanoma cells. For further discussion of the specificity and identity of the cytolytic effector cells generated in assays such as those shown in Table 11, see Gately et al., J. Immunol. 136: 1274-1282, 1986.

Our results demonstrate that purified human CLMF by itself caused proliferation of activated human T lymphocytes, enhanced the cytolytic activity of human NK cells, and augmented human CTL responses. These activities of CLMF, which are similar to those of IL-2, suggest that CLMF, like IL-2, should have immunoenhancing and antitumor effects when used as a single therapeutic agent in vivo. Clearly, CLMF may also have utility in stimulating the growth in vitro of NK/LAK cells and of activated T cells, such as may be derived from tumor infiltrating lymphocytes [S.L. Topalian et al., J. Immunol. 142: 3714-3725, 1989]. In addition, purified CLMF synergized with low concentrations of rIL-2 in causing the generation of human LAK cells in culture and acted additively or synergistically with rIL-2 in facilitating specific CTL responses in vitro. These results suggest that the use of CLMF in combination with rIL-2 might constitute a more optimal antitumor therapy.

TABLE 8

Purified Human CLMF Stimulates the Proliferation of
5 Human PHA-Activated Lymphoblasts

Expt.	Cytokine Added:		³ H-Thymidine Incorporated by PHA-Activated Lymphoblasts (mean cpm \pm 1 S.E.M.)
	Human CLMF ^c (u/ml)	Human rIL-2 (u/ml)	
10			
1a	0	0	10,607 \pm 596
	500	0	70,058 \pm 1,630
	100	0	60,377 \pm 1,927
	20	0	36,018 \pm 321
15	4	0	24,996 \pm 669
	0.8	0	17,765 \pm 790
2b	0	0	9,976 \pm 374
	200	0	60,980 \pm 1,713
20	50	0	38,817 \pm 884
	12.5	0	18,885 \pm 2,132
	3.1	0	13,648 \pm 731
	0	16	80,041 \pm 5,835
25	0	4	21,282 \pm 1,145
	0	1	11,241 \pm 898
	50	4	62,050 \pm 2,408
	12.5	4	40,628 \pm 2,196
30	3.1	4	31,144 \pm 3,754

- a ~~AB~~ cultures in experiment 1 contained goat anti-human rIL-2.
- 35 b No cultures in experiment 2 contained goat anti-human rIL-2.
- c Purified human CLMF from Mono Q FPLC.

TABLE 9

Purified Human CLMF Synergizes with Human rIL-2
in the Generation of Lymphokine-Activated Killer (LAK) Cells
5 in 4-Day Cultures

	Cytokine Added:		% Specific	⁵¹ Cr Release ^a from:
	Human CLMF ^b (u/ml)	Human rIL-2 (u/ml)	K562	Raji
10	0	0	3 ± 1.7	1 ± 0.5
	800	0	7 ± 0.3	1 ± 0.1
	200	0	5 ± 1.1	1 ± 0.4
	50	0	4 ± 3.0	0 ± 0.9
15	0	5	10 ± 2.4	2 ± 0.8
	800	5	41 ± 4.0	11 ± 0.8
	200	5	42 ± 1.9	11 ± 0.3
	50	5	36 ± 2.7	9 ± 0.8
20	12.5	5	28 ± 2.1	7 ± 0.7
	3.1	5	19 ± 0.8	5 ± 0.3
	0.8	5	14 ± 1.2	3 ± 0.8

25 ^a Values represent the means ± 1 S.E.M. of quadruplicate determinations. The spontaneous ⁵¹Cr release values for K562 and Raji were 16% and 14%, respectively.

30 ^b Purified human CLMF from Mono:Q FPLC.

TABLE 10

Purified Human CLMF Causes Activation of Natural
5 Killer (NK) Cells in Overnight Cultures

Cytokine Added:		% Specific ^{51}Cr Release ^a from Raji Cells at Effector/Target Ratio=:		
10	Human CLMF ^b (u/ml)	Human rIL-2 (u/ml)	20/1	5/1
	0	0	10 \pm 0.6	5 \pm 0.4
	40	0	31 \pm 0.4	14 \pm 0.5
	8	0	23 \pm 2.1	12 \pm 0.4
15	1.6	0	15 \pm 0.3	10 \pm 0.6
	0.3	0	12 \pm 1.2	9 \pm 0.2
	0	1	13 \pm 0.4	6 \pm 0.5
	40	1	33 \pm 2.0	17 \pm 0.5
20	8	1	26 \pm 0.8	13 \pm 1.9
	1.6	1	19 \pm 1.1	11 \pm 2.1
	0.3	1	16 \pm 1.0	10 \pm 1.5
	0	5	20 \pm 1.3	13 \pm 0.6
25	40	5	23 \pm 2.0	12 \pm 1.5
	8	5	29 \pm 1.1	16 \pm 0.7
	1.6	5	27 \pm 1.2	13 \pm 0.8
	0.3	5	24 \pm 1.8	13 \pm 1.2
30	0	25	38 \pm 1.4	19 \pm 0.7

a: Each value represents the mean \pm 1 S.E.M. of quadruplicate determinations. The spontaneous ^{51}Cr release was 9%.

35 b: Purified human CLMF from Mono Q FPLC.

**PURIFIED HUMAN CTLs ENHANCES SPECIFIC HUMAN CYTOLYTIC T LYMPHOCYTE
RESPONSES TO ALLOGENEIC MELANOMA CELLS IN VITRO**

TABLE 11

Contents of Cultures:		Hydrocortisone (M)	Melanoma Cells ^a	CTL-2 (U/ml)	CTL ^b (U/ml)	Specific ⁵¹ Cr Release from 2			
Lymphocytes (all fraction ^b)						Expt. 1		Expt. 2	
						HT144	K562	HT144	K562
4	10 ⁻⁴					6 ± 3	-4 ± 1	-2 ± 2	3 ± 4
4	10 ⁻⁴			7.5	10	-3 ± 1	-1 ± 1	-5 ± 1	4 ± 1
4	10 ⁻⁴	HT ^{uv}				0 ± 2	-3 ± 1	-2 ± 2	2 ± 1
4	10 ⁻⁴	HT ^{uv}			10	7 ± 2	-1 ± 1	13 ± 3	-4 ± 1
4	10 ⁻⁴	HT ^{uv}			7.5	5 ± 1	3 ± 1	1 ± 3	3 ± 3
4	10 ⁻⁴	HT ^{uv}			7.5	20 ± 3	10 ± 1	27 ± 9	-6 ± 1
4	10 ⁻⁵	HT ^{uv}				17 ± 3	-4 ± 1	10 ± 1	3 ± 1
4	10 ⁻⁵	HT ^{uv}			10	39 ± 3	-2 ± 2	33 ± 6	5 ± 3
1 + 2				15		33 ± 4	67 ± 3	19 ± 1	47 ± 1

^a In both experiments the contents of duplicate cultures were pooled, washed, resuspended in 1.2 ml TCM, and assayed for lytic activity undiluted and at 1:5 dilution. In experiment 1, the data shown were obtained using the 1:5 dilution of lymphocytes in the cytolytic assay, corresponding to a lymphocyte: target ratio of approximately 4:1. In experiment 2, significant lysis was only when lymphocytes were added undiluted to the lytic assay, and these data are shown in the table. The spontaneous release from HT144 cells was 25% and 31% in experiments 1 and 2, respectively, and for K562 was 18% and 27% in experiments 1 and 2, respectively.

^b Percoll Fraction 4 contained high density lymphocytes recovered from the interface between the 45% and 56% Percoll layers, whereas Percoll fraction 1 + 2 contained lower density lymphocytes harvested from the interfaces between the 35% and 38% and between the 38% and 41% Percoll layers. Percoll fraction 4 contained CTL precursors but few LAK precursors; on the other hand, fraction 1 + 2 was rich in LAK cell precursors.

^c HT^{uv} and HT^y represent HT144 melanoma cells which had been uv-irradiated or gamma-irradiated, respectively.

Example 10

5 CLONING OF A cDNA CODING FOR THE 40 kDa SUBUNIT OF HUMAN CLMF

1) Cell culture and isolation of polyA+ RNA

10 NC 37 cells (subclone 98) were grown in roller bottles as described above and induced with PMA and calcium ionophore for 15.5 hours. The cells were harvested, resulting in a frozen cell pellet of 1.11 grams, 5.25×10^8 cells. A portion of the culture was continued for 3 days, at which point the bioassay titer for CLMF
15 activity read 2200 units/ml, indicating that the cells harvested for isolation of RNA had indeed produced the CLMF activity. Total RNA was isolated from the frozen cells by standard procedures and polyA+ RNA was obtained by affinity chromatography. The yield of polyA+ RNA was 2.5% (w/w) relative to the total amount of RNA
20 input.

2) Establishment of a cDNA library

2 μ g of the above polyA+ RNA were reverse transcribed into
25 cDNA using 150 ng random hexamers as primers. A library in lambda gt10 was established, and 1.5×10^5 clones were amplified for the screening.

30 Use of PCR to Generate a DNA Probe Specific for the 40 KDa CLMF Subunit cDNA

The partial N-terminal sequence of the purified 40 kDa protein is IWELKKDVYVVVELDWYPDAP.... Two primers for use in mixed
primer PCR were designed and synthesized by standard procedures.
35 The forward primer was designed as the coding strand corresponding to amino acids ELKKD in the above sequence, containing all possible

codons for that sequence and having an extension at its 5' end including an EcoRI site and three additional bases adding stability. The sequence of the forward primer is thus 5' ctc gaa ttc gaa/g c/ttn aaa/g aaa/g ga, i.e. a 23mer with 64 different sequences. The reverse primer was designed in the same manner, to represent the antisense strand corresponding to the amino acid sequence YPDAP in the partial N-terminal 40 kDa sequence. The reverse primer thus has the sequence 5' ctc gaa ttc ngg ngc a/gtc ngg a/gta and is a 24 mer containing 256 different sequences. The symbol n stands for any one of the four possible bases a,g,c or t. The primers thus define an amplicon of 72 basepairs in length. After cutting with EcoRI for generating cohesive ends for subcloning, the amplicon size drops to 64 basepairs. Single-stranded cDNA was generated for use in the PCR as described in section 2 above, using polyA+ RNA from induced and, as a control, uninduced cells. 40 ng of either one of those cDNAs were amplified with forward and reverse primers in 100 µl of 10 mM Tris-HCl pH 8.3/50mM KCl/1.5 mM MgCl₂/0.01 % gelatine/200 uM each of the four nucleotides/10 units Taq-polymerase/250 pmoles of each primer. The PCR parameters were as follows: initial denaturation was at 95°C for 7 minutes. Low stringency annealing was performed by cooling to 37°C over 2 minutes, incubating 2 minutes at 37°C, heating to 72°C over 2.5 minutes, extending at 72°C for 1.5 minutes, heating to 95°C over 1 minute and denaturing at 95°C for 1 minute; this low stringency annealing cycle was repeated once. Afterwards, 30 standard cycles were run as follows: 95°C for 1 minute, 55°C for 2 minutes, 72° C for 2 minutes. A final extension was performed at 72°C for 10 minutes. 10% of the total samples were run on a 4% agarose gel, stained and analyzed. The amplicon of the expected size was only detectable in the sample where induced cDNA had been amplified. The remainder of the sample was extracted with phenol, concentrated by precipitation with ethanol and redissolved in 42 µl of water. The sample was digested with 60 units of the restriction enzyme EcoRI in 50 µl at 37°C for 2 hours. The sample was subsequently run on a 6% polyacrylamide gel and the 64 bp amplicon was cut out of the gel and eluted by

- standard procedures. The DNA amplicon was subcloned into the EcoR1 site of the bluescript SK+ plasmid by standard procedures (5). Colonies obtained from the transformation of the E.coli strain DH5 alpha were picked and analyzed for the presence of the 64 bp insert.
- 5 Two positive candidates were sequenced to determine the sequence of the cloned amplicon. It is clear from this analysis that the correct fragment was amplified, since the deduced amino acid sequence matches exactly the partial amino terminal amino acid sequence from the purified 40 kDa protein. This information was
- 10 subsequently used to design a 54 bp long oligonucleotide probe that could be used for screening of the cDNA library. Two oligos were designed, with the following sequence: 5' gag cta aag aaa gat gtt tat gtc gta gaa ttc gat and 5' agg ggc atc cgg ata cca atc caa ttc tac gac ata. These two oligos are partially complementary to form the following
- 15 structure:

5' gagctaaagaagatgtttatgtcgtagaattggat 3'
3' atacagcatcttaacctaaccataggcctacgggga 5'

- 20 Such a structure can be labelled by using klenow fragment and labelled nucleotides such that a high specific activity probe results for the screening of cDNA libraries.

25 4) Screening of cDNA Libraries

- A total of 3×10^5 clones from the amplified library were screened on 6 duplicate filters under the following conditions: 50 ml of 5xSSC/10x Denhardt's/100 µg/ml denatured calf thymus DNA/20% formamide/0.1% SDS/1.5 x 10⁶ cpm of labelled 54 mer at 37°C for 16
- 30 hours. The filters were subsequently washed in 2xSSC at 42°C for 30 minutes, dried and exposed to X-ray film. After overnight exposure with an intensifying screen, 16 possible positives were picked and further analyzed by a second screening round. 10 rehybridizing phage were isolated and their DNA prepared. 8 of those 10 isolates
- 35 looked identical, upon EcoR1 cutting releasing two fragments of .8kb

and .6kb length, indicating a possible internal EcoRI site. Upon blotting and hybridization with the screening probe, only the .6kb fragment showed hybridization. The two fragments were subcloned separately into the EcoRI site of the bluescript SK+ plasmid as described above and were completely sequenced. This analysis showed that both fragments align in one contiguous cDNA of about 1.4kb in length with a naturally occurring internal EcoRI site, since both fragments upon translation showed the presence of reading frames coding for tryptic peptides that had actually been isolated from purified 40Kd protein. The complete sequence of the 40 kDa subunit as deduced from the cDNA is shown in figure 25. The cDNA codes for one open reading frame of 328 amino acids. The protein starts with the initiating Met, followed by another 21 amino acids that make up a classical hydrophobic signal peptide. The N-terminus of mature purified 40 kDa subunit, i.e. IWELKKD..., follows immediately after the signal sequence. The mature protein thus consists of 306 amino acids. The deduced protein sequence contains 4 possible N-linked glycosylation sites, two of which are present in isolated and sequenced tryptic peptides. One of these two sites is used in vivo for the attachment of a carbohydrate side chain. The molecular weight of the mature unglycosylated protein is 34699, the pI is 5.24. The corresponding mRNA is 2.4kb in length and is detectable in a northern blot in steady state RNA only from induced cells.

25

Example 11

Cloning of a cDNA coding for the 35 kDa subunit of human CLMF

30 **Cell culture, isolation of mRNA and establishment of a cDNA library were as described earlier for the cloning of the 40 kDa subunit.**

Use of mixed primer PCR to generate a DNA probe specific for the 35 kDa subunit cDNA

- The partial N-terminal sequence of the purified 35 kDa subunit is ?NLPVATPDPGMF?LHHSQNLLRAV.... Two primers for use in mixed primer PCR were generated by standard procedures. The forward primer was designed as the coding strand corresponding to the amino acids DPGMF in the above sequence, containing all possible codons for that sequence and having an extension at its 5' end including an EcoRI site and three additional bases adding stability. The sequence of this forward primer was thus 5' CTC GAA TTC GAT/C CCN GGN ATG TT -3', i.e. a 23 mer with 32 different sequences. The reverse primer was designed in the same manner, to represent the antisense strand corresponding to the amino acids NLLRA in the partial N-terminal sequence. The reverse primer had the sequence 5' CTC GAA TTC NGC NCG/T NAA/G NAA/G A/GTT, i.e a 24mer with 4096 different sequences. In both primer sequences, N stands for all 4 bases. The two primers thus defined an amplicon 69 bases long. After cutting with EcoRI for generating cohesive ends for subcloning, the amplicon size drops to 61 bases. About 3 µg. of human genomic DNA were amplified with forward and reverse primers in 50 µl of 10 mM Tris-HCl pH 8.3 / 50 mM KCl / 1.5 mM MgCl₂ / 0.01% gelatine / 200 µM each of the four nucleotides / 2.5 units of Taq polymerase / 64 pmoles of forward and 2048 pmoles of reverse primer (to compensate for the greatly differing complexities of the two primers). The PCR cycling parameters were as follows: initial denaturation was at 95° C for 7 minutes. Low stringency annealing was performed by cooling to 37° C over 2 minutes, incubating at 37° C for 2 minutes, heating to 72° C over 2.5 minutes, extending at 72° C for 1.5 minutes, heating to 95° C over 1 minute and denaturing at 95° C for 1 minute; this low stringency annealing cycle was repeated once. Afterwards, 40 standard cycles were run as follows: 95° C for 1 minute, 55° C for 2 minutes and 72° C for 3 minutes. A final extension was performed at 72° C for 10 minutes. About 20% of the samples were run on a 6% polyacrylamide gel and an amplicon of the expected size was detected after staining the gel

- with ethidium bromide. The remainder of the sample was extracted with phenol, concentrated by precipitation with ethanol and redissolved in 17 μ l of water. The sample was digested with 20 units of EcoRI enzyme in 20 μ l for 60 minutes at 37°C. The sample was
- 5 subsequently fractionated on an 8% polyacrylamide gel and the 61 basepair amplicon was cut out of the gel and eluted by standard procedures. The DNA amplicon was subcloned into the EcoRI site of the Bluescript plasmid SK+ by standard procedures. Colonies obtained from the transformation of the strain DH5 alpha were analyzed for
- 10 the presence of the 61 basepair insert. Two candidates were sequenced to determine the sequence of the subcloned amplicon. One of the two clones contained the correct sequence, since translation of that sequence resulted in the amino acid sequence expected from the purified protein. Based on this information, two synthetic
- 15 oligonucleotides were designed, with the following sequences:

5'gatccgggaatgttcccatgccttcaccactccc 3'

3'gtacggaagtgtgagggttttgaggatgcccca 5'

- 20 Such a structure can be labelled using radiolabelled nucleotides with the Klenow fragment to very high specific activities for library screening.

Screening of a cDNA library:

- 25 A total of 10^6 clones from the amplified 16 h library were screened on 40 duplicate filters with the above probe under the following conditions: 400 ml of 5X SSC / 20% formamide / 10 X Denhardt's / 100 μ g/ml denatured calf thymus DNA / 0.1 % SDS / 3.8
- 30 $\times 10^7$ cpm labelled probe at 37 ° C overnight. The filters were subsequently washed in 2X SSC at 40° C and exposed to X-ray film with a screen overnight. Six potential positives were picked from this first round of screening and analyzed by a second round of plaque hybridization, as above. One clone was picked for the final
- 35 analysis. Upon preparing phage DNA, the clone was found to contain

two EcoR1 fragments of about 0.8kb and 0.3 kb in size. The two fragments were subcloned separately into the Bluescript SK+ plasmid and sequenced. This analysis showed that the two fragments align into one contiguous sequence of about 1.1kb total length with a naturally occurring internal EcoR1 site. The complete sequence of the cDNA and the deduced amino acid sequence for the 35 kDa CLMF subunit are shown in figure 26. The cDNA codes for an open reading frame of 219 amino acids, starting with the initiating Met at position 1. The following 21 amino acids constitute a classical hydrophobic signal sequence. Immediately following the signal peptide, the N-terminus of the mature 35 kDa protein starts with the sequence RNLPVAT.... Purified 35 kDa protein had yielded the sequence ?NLPVAT.... The mature 35 kDa protein thus consists of 197 amino acids, containing three possible N-linked glycosylation sites and 7 cys-residues. The molecular weight of mature unglycosylated protein is 22513, and the pI is 6.09. The corresponding mRNA is 1.4Kb in length and is only detectable in RNA from cells that had been induced for CLMF for at least 6 hours.

Example 12

Expression of biologically active recombinant CLMF in COS-cells

The two subunits for CLMF were engineered for expression in mammalian cells as follows:

40 kDa subunit

The two EcoR1 fragments constituting the full length cDNA for the 40 kDa CLMF subunit were ligated to an expression vector similar to pBC12 [See B. Cullen, Meth. Enzymology 152, 684,703, (1987)], except that the cDNA expression is driven off the SV40 early promoter/enhancer. Clones containing the two inserts in the proper orientation to each other were selected by colony hybridization with a synthetic oligonucleotide that spans the internal EcoR1 site in the 40 kDa cDNA. This oligonucleotide has the following sequence:

5' CTG AAG CCA TTA AAG AAT TCT CGG CAG GTG 3'. It was labelled by kinasing using standard procedures. Clones were subsequently analyzed for proper orientation of insert to vector by the polymerase chain reaction procedure, using as forward primer a primer specific for sequences in the SV 40 early promoter and as reverse primer an oligonucleotide corresponding to the 40 kDa cDNA sequence positions no. 851-868. Clones with the correct orientation will give a PCR amplicon of 885 bp. Eight out of 20 clones tested gave the predicted fragment, and one was chosen for further study.

35 kDa subunit

The full length cDNA for the 35 kDa subunit was amplified out of the original lambda phage by PCR, using primers situated to the left and right of the EcoRI site in lambda gt10 (primers were New England Biolab Articles No. 1231 and 1232). The resulting PCR amplicon was blunt-end ligated into the EcoRV site of the bluescript plasmid SK+ and the DNA propagated. DNA sequencing showed that the orientation of the cDNA insert within the plasmid was such that the end of the cDNA corresponding to the 5' end of the mRNA was close to the ClaI site in the polylinker. The insert was thus released by cutting with ClaI, filling out this end with T4 DNA polymerase and cutting secondarily with Not I. The resulting fragment was gel-purified and subcloned into an expression plasmid based on the bluescript vector and containing the SV40 early promoter to drive expression of inserted cDNAs. The sites in the expression plasmid used were a blunt-ended PstI site at the 5' end and a Not I site at the 3' end of the cDNA. One clone was chosen for further study after ascertaining its structure by PCR as above for the 40 kDa construct.

Expression of the two cDNAs in COS-cells

The DNAs for the expression constructs of the 40 kDa and 35 kDa subunits were introduced into COS cells on 6 cm diameter plates by the DEAE Dextran transfection procedure (7×10^5 cells/dish plated; 1 μ g DNA per dish). 24 hours after the transfection, the cells

were fed with standard tissue-culture medium containing 1% Nutridoma instead of the fetal bovine serum and the supernatants were collected after 40 hours and filtered through a 0.45 μ filter.

- 5 Supernatant fluids from cultures of COS cells which had been transfected with cDNA encoding the 35 kDa CLMF subunit or the 40 kDa CLMF subunit or with both cDNAs were tested for CLMF activity in the T cell growth factor assay (Table 12). As shown in the table, COS cells which had been transfected with only one of the subunit
10 cDNAs did not release biologically active CLMF into the culture fluid. However, COS cells which had been transfected with both subunit cDNAs produced biologically active CLMF. By comparing the amount of lymphoblast proliferation induced by the culture fluid from doubly transfected COS cells to the amount of proliferation induced
15 by purified NC-37-derived CLMF, the concentration of CLMF activity in the culture fluid was estimated to be 374 units/ml. Assuming a specific activity of 8×10^7 units/mg CLMF protein, this result suggests that the fluid from cultures of doubly transfected COS cells contained approximately 4.7 ng/ml of recombinant CLMF.
20

TABLE 12

T Cell Growth Factor Activity of Recombinant CLMF
Expressed in COS cells

Cytokine added:	Concentration:	³ H-Thymidine Incorporated by PHA-Activated Lymphoblasts (mean cpm ± 1 S.E.M.)
None		14,587 ± 363
Natural CLMF*	200 units/ml	79,846 ± 854
Natural CLMF	40 units/ml	59,093 ± 2039
Natural CLMF	8 units/ml	39,180 ± 545
Natural CLMF	1.6 units/ml	25,996 ± 763
Supernatant fluid from cultures of COS cells transfected with:		
35 Kda CLMF subunit cDNA	1/5 dilution	15,332 ± 797
35 Kda CLMF subunit cDNA	1/25 dilution	12,149 ± 379
40 Kda CLMF subunit cDNA	1/5 dilution	14,883 ± 1039
40 Kda CLMF subunit cDNA	1/25 dilution	13,689 ± 110
35 Kda + 40 Kda CLMF subunit cDNAs	1/5 dilution	66,228 ± 166
35 Kda + 40 Kda CLMF subunit cDNAs	1/25 dilution	47,673 ± 275
Mock transfected	1/5 dilution	14,366 ± 628
Mock transfected	1/25 dilution	14,426 ± 173

*Purified NC-37-derived CLMF from Mono Q FPLC.

Example 13

Preparation, Characterization and Purification of Hybridoma 5 Antibodies

Lewis rats (Charles River Laboratories, Wilmington, Ma) were initially immunized by the intraperitoneal route (i.p.) with partially purified CLMF mixed with an equal volume of Freund's complete
10 adjuvant (Gibco). The rats were injected i.p. with booster immunization of CLMF mixed with Freund's incomplete adjuvant (Gibco) according to the schedule in Table 13. For preparation of activated spleen cells, one rat was injected i.v. with partially purified CLMF on two successive days starting 4 days prior to the cell fusion
15 (Table 14). Spleen cells were isolated from this rat and fused with NSO cells at a ratio of 1:1 (spleen cells:NSO cells) with 35% polyethylene glycol (PEG 4000, E. Merck). The fused cells were plated at a density of 5×10^4 cells/well/ml in 48 well plates in IMDM supplemented with 15% FBS, glutamine (2 mM), beta-
20 mercaptoethanol (0.1 mM), gentamycin (50 ug/ml), HEPES (10 mM) and 15% P388D1 cell supernatant. Hybridoma supernatants were screened for specific CLMF antibodies in 4 assays: 1) immunoprecipitation of ^{125}I -labelled CLMF, 2) immunodepletion of CLMF bioactivity, 3) western blotting with CLMF and 4) inhibition of
25 ^{125}I -CLMF binding to PHA-activated PBL blast cells. Hybridoma cell lines secreting anti-CLMF antibodies were cloned by limiting dilution. Antibodies were purified from large scale hybridoma cultures or ascites fluids by affinity chromatography on protein G bound to cross-linked agarose according to the manufacturer's protocol
30 (Gammabind G, Genex, Gaithersburg, MD).

Isolation and Identification of Monoclonal Antibodies Specific
for CLMF. Serum isolated at the 3rd bleed from the rat immunized with partially purified CLMF (Table 13) neutralized CLMF bioactivity
35 (5 units/ml) as determined in the TGF assay (Fig. 27). This neutralization could be blocked by adding excess CLMF (200

- units/ml) demonstrating that the neutralization by the antiserum was specific for CLMF (Fig. 27). Normal rat serum did not neutralize CLMF bioactivity (Fig. 27). Spleen cells isolated from this rat were fused with NSO cells and the resulting hybridomas were initially screened for CLMF-specific antibodies by immunoprecipitation of ¹²⁵I-labelled CLMF.

- The radioiodinated partially purified CLMF preparation contains predominantly the CLMF 75 kDa heterodimer, a small amount of the free CLMF 40 kDa subunit and two other proteins of approximately 92 kDa and 25 kDa (Fig. 28). The ¹²⁵I-labelled CLMF preparation retained CLMF bioactivity in the TGF assay, indicating that the labelling procedure did not significantly alter the configuration of the CLMF molecule. The CLMF immunized rat serum immunoprecipitated the 75 kDa heterodimer and the free 40 kDa subunit (Lanes 6 and 8, Fig. 28) whereas normal rat serum did not immunoprecipitate these radiolabelled proteins (Lanes 7 and 9, Fig. 28). Four individual monoclonal antibodies also immunoprecipitated the 75 kDa heterodimer and the free 40 kDa subunit (Fig. 28) but did not immunoprecipitate the 92 kDa or 25 kDa labelled proteins. The immunoprecipitation assay identified twenty hybridomas which secreted anti-CLMF antibodies (Table 14). All the antibodies immunoprecipitated the radiolabelled 75 kDa heterodimer and the free 40 kDa subunit as determined by SDS/PAGE and autoradiography (data shown for 4 representative antibodies in Fig. 28).

- After initially identifying specific CLMF antibodies in the immunoprecipitation assay, the antibodies were tested for their ability to immunodeplete CLMF bioactivity as assessed by the TGF and LAK cell induction assays. Increasing amounts of CLMF cause a dose dependent increase in the proliferation of PBL blasts in the TGF assay as measured by the incorporation of ³H-thymidine into the dividing blast cells (Fig. 29). Immunodepletion of CLMF activity by immobilized anti-CLMF antibodies occurs in a dose dependent manner (Fig. 29). Aliquots (0.4 and 0.1 ml) of hybridoma

- supernatant solution will completely deplete 50 and 200 units/ml of CLMF activity from the culture medium. 0.025 ml of supernatant solution will completely deplete 50 units/ml but only approximately 50% of 200 units/ml. 0.0062 ml of hybridoma supernatant shows
- 5 even less depletion of 50 and 200 units/ml of CLMF. An aliquot (0.4 ml) of an anti-IL-1 receptor antibody supernatant solution shows no immunodepletion of CLMF bioactivity.

- Increasing amounts of CLMF also cause a dose dependent
- 10 increase in the lysis of target cells by LAK cells as measured by the release of ^{51}Cr in the LAK cell induction microassay (Fig. 30). The immobilized anti-CLMF antibodies also deplete in a dose dependent manner the CLMF activity in the LAK cell induction assay (Fig. 30). These data confirm that the antibodies which immunoprecipitate the
- 15 75 kDa labelled protein from the radiolabelled partially purified CLMF preparation are specific for CLMF. The data also demonstrate that the radiolabelled 75 kDa protein is the protein responsible for CLMF bioactivity.

- 20 Identification of the CLMF Subunit Bound by the Monoclonal Antibodies. CLMF is a 75 kDa heterodimer protein composed of 40 kDa and 35 kDa subunits. Western blot analysis was used to determine if the monoclonal anti-CLMF antibodies recognized the 40 kDa or the 35 kDa subunits. Highly purified 75 kDa CLMF
- 25 heterodimer was separated by non-reducing SDS/PAGE and transferred to nitrocellulose membrane (Fig. 31). In addition, purified CLMF, which was composed of approximately 95% free 40 kDa subunit and 5% 75 kDa heterodimer, was separated by both non-reducing and reducing SDS/PAGE and the proteins were transferred
- 30 to nitrocellulose membrane (Fig. 32). Individual nitrocellulose strips containing the non-reduced 75 kDa CLMF heterodimer (Fig. 31), the non-reduced 40 kDa subunit (top panel Fig. 32) and the reduced 40 kDa subunit (bottom panel Fig. 32) were probed with monoclonal anti-CLMF antibodies, control monoclonal antibody, rat anti-CLMF
- 35 serum and control rat serum. The monoclonal anti-CLMF and rat polyclonal anti-CLMF antibodies bind specifically to an

approximately 75 kDa heterodimer on the strips containing non-reduced 75 kDa CLMF while the control antibody preparations do not show this binding activity (Fig. 31). All the monoclonal and rat polyclonal anti-CLMF antibodies recognize the non-reduced 40 kDa subunit (top panel, Fig. 32). However, only the rat polyclonal antiserum and three monoclonal antibodies, 8E3, 9F5 and 22E7, bind to reduced 40 kDa subunit protein (bottom panel, Fig. 32). These data demonstrated that all the monoclonal antibodies were specific for the 40 kDa subunit of CLMF.

Identification of a CLMF Receptor on PHA-Activated

Lymphoblasts. The previous data demonstrated that the monoclonal anti-CLMF antibodies immunoprecipitated ^{125}I -labelled CLMF, immunodepleted CLMF bioactivity and bound to the 40 kDa subunit of CLMF. However, the antibodies present in the hybridoma supernatant solutions could not be directly tested for their ability to neutralize CLMF bioactivity in the TGF or LAK cell induction assays due to non-specific inhibitory effects of supernatant solutions containing control antibodies. Our previous work with IL-2 monoclonal antibodies demonstrated that antibodies which would block ^{125}I -IL-2 binding to IL-2 receptor bearing cells would also neutralize IL-2 bioactivity. Since receptor binding assays are usually unaffected by addition of hybridoma supernatant solutions or other substances, a CLMF receptor binding assay was developed to evaluate the anti-CLMF antibodies for inhibitory/neutralization activity. A CLMF receptor binding assay was configured with ^{125}I -labelled CLMF and the PHA-activated peripheral blood lymphoblasts (Fig. 33). The binding of ^{125}I -CLMF to the PHA-activated lymphoblasts was saturable and specific (Fig. 33). Scatchard plot analysis [See Scatchard, G. Ann. N.Y. Acad. Sci. 51, 660-672 (1949)] of the equilibrium binding data indicated that the apparent dissociation constant for ^{125}I -CLMF binding to the receptor is approximately 200 pM and that each lymphoblast has about 700-800 receptors. Since the serum from the rat immunized with CLMF showed neutralization of CLMF bioactivity, it was tested for inhibition of ^{125}I -CLMF binding to the lymphoblasts (Fig. 34). The

rat immune serum blocks 50% of ^{125}I -labelled CLMF binding at approximately a 1/500 dilution, while the control rat serum does not show any inhibition at this dilution. With the specificity of the receptor binding assay established, hybridoma supernatant solutions
5 were tested for antibodies which would inhibit ^{125}I -CLMF binding to lymphoblasts.

The degree of inhibition of ^{125}I -CLMF binding to the lymphoblasts was determined at a 1/2 dilution of each hybridoma
10 supernatant solution (Fig. 35). Twelve hybridoma supernatant solutions inhibited by greater than 60% ^{125}I -CLMF binding to the lymphoblasts. The antibodies present in these supernatant solutions have been classified as inhibitory/neutralizing antibodies. Six hybridoma supernatant solutions inhibited ^{125}I -labelled CLMF
15 binding by less than 40% and were classified as non-inhibitory/non-neutralizing antibodies. Control antibody inhibited by approximately 10% the ^{125}I -CLMF binding to the lymphoblasts.

Three inhibitory antibodies, 7B2, 2A3 and 4A1, and two non-inhibitory antibodies, 6A3 and 8E3, were purified from ascites fluid
20 by protein G affinity chromatography on GammaBind G (Genex, Gaithersburg, MD) columns. Antibodies 4A1, 2A3 and 7B2 inhibit in a dose dependent manner ^{125}I -CLMF binding to the lymphoblasts with IC_{50} concentrations of 0.7 $\mu\text{g}/\text{ml}$, 7 $\mu\text{g}/\text{ml}$ and 9.5 $\mu\text{g}/\text{ml}$,
25 respectively (Fig. 36). Antibodies 6A3 and 8E3 do not block ^{125}I -CLMF binding at concentrations of 100 $\mu\text{g}/\text{ml}$ (Fig. 36). These data demonstrated that the original classification of each antibody as either inhibitory or non-inhibitory was correct.

30 Direct Neutralization of CLMF Bioactivity by Antibodies. To determine if the antibodies classified as inhibitory by the CLMF receptor binding assay would directly neutralize CLMF bioactivity, each inhibitory antibody was tested for neutralizing activity in the TGF assay (Table 15). Two inhibitory antibodies, 4A1 and 7B2,
35 demonstrated a dose dependent neutralization of CLMF bioactivity (40 units/ml) from .03 to 100 $\mu\text{g}/\text{ml}$, with IC_{50} concentrations of

approximately 1 $\mu\text{g/ml}$ and 80 $\mu\text{g/ml}$, respectively. These data confirmed that antibodies inhibiting ^{125}I -CLMF binding to the CLMF receptor would also neutralize CLMF bioactivity.

5

TABLE 13

Immunization Schedule:

10	Date	CLMF (10^8 units/mg)	mg	Total Protein (μ g)	Spec. Activity (U/mg)	Purity (%)
	3/28/89	1×10^4	.1 μ g	15	6.7×10^5	6.7
	4/10/89	1.2×10^4	.1 μ g	?	6×10^5	.6
15	5/3/89	1st bleed				
	5/18/89	2.2×10^5	2 μ g	75	2.9×10^6	2.9
20	6/7/89	2nd bleed				
	6/29/89	6.3×10^4	.63 μ g	83	7.5×10^5	.75
	7/21/89	1.2×10^5	1.2 μ g	24	5×10^6	5.0
25	8/2/89	3rd bleed				
	10/19/89	2.1×10^6 (i.v.)				
	10/20/89	2.1×10^6 (i.v.)				
	10/23/89	Fusion				
30						

Table 14: Monoclonal Anti-CLMF Antibodies (40 kDa Subunit Specific)

5	Antibody	Western Blot ¹	125I-CLMF/Receptor Assay	Neutralization
	Red.	N.R.	(% Inhibition) ²	of Bioactivity ³
<u>Inhibitory/Neutralizing</u>				
10	7B2	- ++	95	+
	2A3	- ++	99	+
	1B8	- +/-	60	ND ⁴
	1C1	- ++	81	ND
15	4A1	- +	98	+
	4C8	ND ND	68	ND
	4D1	- +	100	ND
	6A2	- +/-	75	ND
	7A1	+/- ++	94	ND
	8A1	- +	99	ND
20	8A2	- ++	83	ND
	9C8	- +	62	ND
	22E7	++ ++	91	ND
<u>Non-Inhibitory/Non-Neutralizing</u>				
25	8E3	+ ++	35	-
	9F5	+ ++	18	ND
	4A6	- -	17	ND
	6A3	- +	20	-
	8C4	- ++	33	ND
30	8E4	- ++	1	ND
	39B3	ND ND	46	ND
	Control	- -	12	-
35	1. Western blots: N.R. is non-reduced and Red. is reduced SDS/PAGE For the western blots, a CLMF sample containing 5% 75 kDa heterodimer and 95% free 40 kDa subunit were separated on 10% SDS/PAGE and western blots prepared as described in me. The blots were scored as strongly positive (++), positive (+), weakly positive (+/-), negative (-).			
40	2. CLMF receptor binding assay: An antibody was considered inhibitory if it would block more than 60% of radiolabelled CLMF binding to the PHA activated PBL cells			
45	3. Neutralization of CLMF bioactivity as assessed by the TGF assay: An antibody was considered neutralizing if it would block more than 50% proliferation at 20 µg/ml. The results are presented as positive (+) or negative (-).			
	4. ND: Not Determined			

Table 15. Neutralization of CLMF Bioactivity by Monoclonal Anti-CLMF.

5 Assay Contents:

	CLMF ^a	Antibody ^b	Total ³ H-Thymidine Incorporation	% Neutralization ^c
10	none	none	9923 ± 439	
	CLMF	none	25752 ± 592	
	CLMF	4A1		
15		100 µg/ml	12965 ± 938	81
		20	12215 ± 663	86
		4	12985 ± 269	81
		.8	19932 ± 1016	37
		.16	22379 ± 410	21
20		.03	25405 ± 1093	2
	CLMF	7B2		
		200 µg/ml	10763 ± 878	96
		100	15083 ± 406	67
25		20	23690 ± 1228	13
		4	25849 ± 1408	0
	CLMF	Control		
		200 µg/ml	27654 ± 1086	0
30		100	22221 ± 381	22
		20	27335 ± 620	0

35 ^a Purified CLMF was used in the TGF assay at a concentration of 40 units/n

^b Purified antibodies were added at the concentrations indicated in the tab

^c Reduction of ³H-thymidine incorporation to the level seen in the absence added cytokines was considered to be 100% neutralization.

40

Example 14

Preparation of Antibodies Against a Synthetic Peptide Fragment of the 35,000 dalton Subunit of CLMF

5

A peptide, comprising amino acids 3-13 of the NH₂-terminal sequence of the 35 kDa CLMF subunit and a COOH-terminal cysteine (L-P-V-A-T-P-D-P-G-M-F-C), was synthesized by solid-phase peptide methodology, purified by HPLC, and conjugated to keyhole limpet hemocyanin via the methylated bovine serum albumin procedure. Two rabbits were immunized intradermally with the conjugated peptide in Freund's complete adjuvant (300 µg peptide/rabbit). Six weeks after immunization, rabbits were boosted with free peptide (100 µg, intravenously) and KLH-conjugated peptide (150 µg, subcutaneously) dissolved in PBS. Serum samples were prepared from bleedings taken 7 days later. The boosting and bleeding schedule was repeated every 4-5 weeks.

Serum samples from the first and second bleedings from each rabbit were evaluated for reaction with the synthetic peptide in a direct ELISA assay. The synthetic, free peptide was coated on microtiter plates at 4 ng/ml and 20 ng/ml, and the plates were washed and blocked with bovine serum albumin. Serum samples were tested at various dilutions (Table 16), and antibody reactivity was detected with the use of a second antibody (HRP-conjugated goat anti-rabbit IgG) with o-phenylenediamine as substrate. Absorbance values were read at 490 nm after addition of H₂SO₄ to stop the reaction. The results indicate that antibody was produced in both rabbits against 35,000 dalton CLMF peptide (Table 16). In separate experiments, we verified that the antibody was specific for the peptide since (a) serum from non-immunized rabbits does not react with the peptide in ELISA, (b) sera from rabbits immunized with the synthetic peptide do not react with a peptide fragment from the 40,000 dalton CLMF subunit and (c) purified IgG from the serum samples also reacts with the synthetic peptide.

A serum sample from one of the rabbits (first bleed) was tested by Western blot analysis for reactivity with 75 kDa CLMF and with the 35 kDa CLMF subunit (Fig. 37). Partially purified CLMF (approximately 120 µg/ml) was run on SDS-PAGE, transferred to
5 nitrocellulose, and treated with a 1:500 dilution of the rabbit anti-CLMF peptide antiserum. Antibody reactivity was detected by use of biotinylated goat anti-rabbit IgG and alkaline phosphatase-conjugated streptavidin. The anti-CLMF peptide antibody was found to react both with nonreduced 75 kDa CLMF protein and with the
10 reduced 35 kDa CLMF subunit (Fig. 37).

Although the antibodies produced in this example were polyclonal, a similar approach could be used to prepare monoclonal antibodies to the 35 kDa subunit of CLMF. The synthetic peptide
15 used in this example or other synthetic peptides based on the amino acid sequence of the 35 kDa CLMF subunit (Fig. 26) could be used to immunize rats. Fusions could be performed and hybridoma cultures screened for the production of monoclonal anti-CLMF antibodies as described above.

ELISA ASSAY OF SERUM SAMPLES FROM RABBITS IMMUNIZED
AGAINST A KIT-CONJUGATED SYNTHETIC PEPTIDE FROM THE
35,000 DALTON SUBUNIT OF CME

TABLE 16

Rabbit No.	Serum Source	Bleed No.	Free Peptide (ng/ml)	A ₄₉₀ Serum Dilutions (to 10)				
				1x10 ⁴	2.5x10 ⁴	6.2x10 ⁴	1.6x10 ⁵	3.9x10 ⁵
1		1	20	2.7	2.1	1.4	0.8	0.5
1		1	4	0.7	0.4	0.3	0.2	0.1
1		1	0	0.05	0.05	0.05	0.05	0.05
1		2	20	2.5	1.8	1.1	0.6	0.3
1		1	4	0.9	0.6	0.4	0.2	0.1
1		2	0	0.05	0.05	0.05	0.05	0.05
2		1	20	2.7	2.2	1.6	0.9	0.5
2		1	4	0.9	0.5	0.3	0.2	0.1
2		1	0	0.05	0.05	0.05	0.05	0.05
2		2	20	2.2	1.5	1.0	0.5	0.3
2		2	4	0.7	0.4	0.2	0.1	0.1
2		2	0	0.05	0.05	0.05	0.05	0.05

Example 15*

Synthesis of IgE is dependent upon the balance between the production of IL-4 and IFN- γ at the sites of T/B cell interactions (1-3). IL-4 may promote IgE synthesis not only via a direct effect on B cells, by directing the switching to IgE (4-5), but also by regulating the production of other molecules or cytokines involved in IgE regulation. For example, IL-4 markedly inhibits IFN- γ production by human lymphocytes stimulated by mitogen or allogeneic cells (6-7). Similarly, IFN- γ suppresses the in vivo synthesis of IgE not only by antagonizing the effect of IL-4 on the switching to IgE but also by inhibiting the proliferation of IL-4-producing TH2 lymphocytes (9-10) or by directing the differentiation of naive precursor T cells into T cells producing IFN- γ but not IL-4 (11). Lymphokines other than IL-4 and IFN- γ may also have an important role in the regulation of IgE synthesis. Interferon- α , a cytokine mainly produced by accessory cells, is a potent inhibitor of the in vitro and in vivo synthesis of mouse and human IgE (13). IFN- α also counteracts the effect of IL-4 on the switching to IgE and most interestingly, like IFN- γ , it inhibits the in vivo production of IL-4 and enhances that of IFN- γ (12). Interleukin-12 is a novel cytokine, which like IFN- γ and IFN- α , may be involved in protective immunity against infectious agents such as viruses. Also known as NKSF (Natural Killer Cell Stimulatory Factor) or as CLMF (Cytotoxic Lymphocyte Maturation Factor) IL-12 is a 75 kDa heterodimeric glycoprotein displaying several in vitro activities including: (a) the enhancement, in synergy with IL-2, of the maturation of cytotoxic T cells and of LAK cells (lymphocyte activated killer cells); (b) the increase of the cytotoxic activity of NK cells (15); (c) the promotion of the proliferation of activated T cells and NK cells (16) and; (d) the induction of IFN- γ production by resting or activated peripheral blood NK cells and T cells (17). IL-12 is a strong inhibitor of the T cell-dependent synthesis of IgE by IL-4-stimulated peripheral blood mononuclear cells and the IgE inhibition may be observed in the absence of IFN- γ production.

35

* Provided by Dr. Guy Delespesse, University of Montreal.

Reagents

- Human rIL-4 was obtained from CIBA-GEIGY, Basle, Switzerland (Dr. H. Hofstetter), anti-CD40mAb 89 (18) was received from (Schering Plough, Dardilly, France (Dr. J. Banchereau)); hydrocortisone was obtained from Sigma (St. Louis, MO); PWM was from Gibco Laboratories (Grand Island, NY); anti-IFN- γ neutralizing mAb was purchased from Genzyme, (No. 1598-00 Boston, MA). In preliminary titration experiments, this antibody (25 μ g/ml) completely neutralized the suppressive activity of 500 IU/ml of IFN- γ on the IL-4-stimulated synthesis of IgE by PBMC. IgE (ng/ml) in IL-4-stimulated cultures was 30 ± 4 as compared to 9.8 ± 2 in the presence of IFN- γ (500 IU/ml) and to 31.7 ± 3.8 in the presence of both IFN- γ and anti-IFN- γ mAb. Anti-Lolp1 mAb is a mouse IgG1 antibody directed against the pollen antigen Lolp1 (20).

Human rIL-12 and antibodies to IL-12

- Human rIL-12 was produced by cotransfection of COS cells with a 1:1 molar ratio of the two subunit cDNAs of IL-12 as described by Gubler et al. (13). Crude supernatant fluid from cultures of doubly transfected cells was used as the source of rIL-12 in these experiments. Supernatant fluid from cultures of mock transfected COS cells was used as a control. Monoclonal anti-IL-12 antibody was a 1:1 mixture of two rat monoclonal anti-human IL-12 antibodies, 4A1 and 20C2, which were isolated and purified as previously described (19). The 4A1 antibody is specific for the 40 kDa subunit of human IL-12, and its isotype is IgG2b. The 20C2 antibody appears to react with the 35 kDa subunit of IL-12, and its isotype is IgG1. These two antibodies were previously found to synergize in blocking IL-12-stimulated proliferation of human PHA-activated lymphoblasts.

Cell preparations and culture conditions

- Cells were prepared and cultured as described (20,21): Briefly, peripheral blood mononuclear cells (PBMC) were isolated from heparinized venous blood of healthy individuals by centrifugation over Ficoll-Metrizoate. Umbilical cord blood was collected in heparin-containing tubes and was sedimented 45 min at 37°C with dextran (10% V/V; mol.wt. 200,000, Baker Chemicals Co., Phillipsburg, NJ); the leukocyte-rich plasma was then layered on Ficoll-Metrizoate. Cells were cultured in HB101 culture medium (Hana Biologics Inc., Alameda, CA) supplemented with 5% fetal calf serum (Flow Labs, McLean, VA) penicillin (100 U/ml), streptomycin (100 µg/ml, L glutamine (2 mM) (Gibco Laboratories), sodium pyruvate (10 mM) and Hepes (10 mM). Cells (2×10^5 in 0.2 ml) were cultured in four replicates or more in round-bottomed 96-well tissue culture plates (Linbro) for 12 days in a humidified atmosphere of 5% CO₂ and 95% air. For the induction of IgE synthesis, cultures were supplemented with IL-4 at the final concentration of 10 ng/ml; this concentration was found to be optimal for the induction of IgE synthesis and the suppression of IFN-γ production in mixed-lymphocyte cultures.

RIAs

- Immunoglobulins were measured in cell-free culture supernatants by means of solid-phase RIAs exactly as described (20,21); IFN-γ was measured by a commercially available RIA (Centocor Co., Malvern, PA) with a sensitivity of 1 IU/ml. The net synthesis of Igs and of IFN-γ was determined by subtracting the values measured in the culture supernatants of cycloheximide-treated cells (50 µg/ml) from those of untreated cells. In preliminary experiments, where the levels of IFN-γ in the supernatants of IL-12 stimulated cultures were determined at days 2, 4, 6 and 8, we found, in agreement with a previous report (17), that a plateau was obtained between day 4 to day 6. IFN-γ was therefore routinely measured at day 6; to this end,

50 µl of culture supernatants were collected and replaced by the same volume of fresh culture medium. All the culture supernatants were stored at -20°C until the assay.

5 Northern blot analysis

- Northern blot analysis was carried out exactly as described (6). Total RNA was extracted from cultured PBMC by the guanidium-thiocyanate method with CsCl gradient modification and quantified by measurement of absorbance at 260 nm. The samples (20 µg per lane) were subjected to electrophoresis in formaldehyde-containing 1% agarose gel, and transferred to nylon membrane (Biotrans, ICN, Irvine, CA). The membrane was baked 2 h at 80°C under vacuum, prehybridized in 50% formamide - 5 X Denhardt's - 5X SSC - 10 mM EDTA - 50 mM sodium phosphate pH 6.8 - 0.1% SDS - 250 µg/ml salmon sperm DNA and incubated overnight at 42°C with ³²P labelled cDNA probe in the same buffer. The probes used for the detection of the germ-line and the mature form of Ce mRNA were described by Jaraba et al. (22). A 0.74 Kb Sma I fragment overlapping the germline exon was used to detect germ-line Ce transcript, and the 0.88 Kb Hinf I fragment encompassing most of the Ce1 exon and the totality of the Ce2 exon was used to detect both the productive and the germ-line Ce mRNAs.

2.5 IL-12 suppresses IgE synthesis

- As seen in table 17, IL-12 (60 pM) significantly suppresses the production of IgE and increases the synthesis of IFN-γ by PBMC cultured in the presence of a saturating concentration of IL4 (10 ng/ml). IL-4 significantly but incompletely suppresses the IL-12-induced production of IFN-γ; and it totally abolishes the spontaneous production of IFN-γ. The effects of IL-12 on IgE and IFN-γ production are dose-dependent and they are completely abolished by neutralizing anti-IL-12 mAbs (Figure 38). The production of IgG, IgA and IgM in IL-4-stimulated cultures is not significantly affected by IL-12. However, IL-4 does not induce the production of IgM, IgA or IgG (with the exception of IgG4) (4). The effect of IL-12 on

pokeweed mitogen (PWM) - induced IgE synthesis was examined. In 3 consecutive experiments, IL-12 (60 pM) had no significant effect on the PWM-induced synthesis of IgG (1.6 ± 0.1 versus 1.3 ± 0.6 $\mu\text{g/ml}$; mean ± 1 SD), IgM (1.2 ± 0.4 versus 1.1 ± 0.6 $\mu\text{g/ml}$) and of IgA (1.9 ± 0.6 versus 2.1 ± 0.7 $\mu\text{g/ml}$).

As shown in Fig. 39, IL-12 strongly suppresses the expression of the mature but not of the germ-line C ϵ transcript. This indicates that IL-12 suppresses the synthesis of IgE and possibly inhibits the switching to IgE.

IL-12 appears to suppress IgE synthesis by a mechanism which is distinct from that of IFN- γ . The effect of IL-12 on the synthesis of IgE by umbilical cord blood mononuclear cells (CBMC) costimulated with IL-4 and hydrocortisone was tested. These cells were selected because of their impaired capacity to produce IFN- γ (23) and because exogenous IFN- γ was found to increase rather than to inhibit their synthesis of IgE following stimulation with IL-4 (6). Hydrocortisone (HC) was added to IL-4 stimulated CBMC for two reasons: (i) HC inhibits the production of IFN- γ , even that induced by IL-12; (ii) HC strongly increases the IL-4 stimulated synthesis of IgE, even in the absence of IFN- γ production (24). As seen (table 18), IL-12 markedly inhibits IgE synthesis by neonatal cells cultured in the presence of IL-4 plus hydrocortisone and producing little or not detectable IFN- γ . Moreover, the suppression is unchanged in the presence of a large excess of neutralizing anti-IFN- γ mAb. Thus, it appears that IL-12 can inhibit IgE synthesis by a mechanism which is independent of IFN- γ .

As shown above, picomolar concentrations of IL-12 markedly inhibit the synthesis of IgE by IL-4-stimulated PBMC. The suppression of IgE is observed at the protein and the mRNA levels and it is completely overridden by neutralizing antibodies to IL-12. Given that the production of IgE by IL-4-stimulated lymphocytes involves the switching of precursor B cells to IgE rather than the selective expansion and differentiation of IgE committed B cells (4,

25), the results suggest that IL-12 inhibits the switching to IgE. Consistent with an isotype-specific activity of IL-12, no influence is found of this lymphokine on the production of the other classes of Ig by IL-4-stimulated or by PWM-stimulated PBMC. The data do not
5 exclude an effect of IL-12 on the production of IgG4, the only human isotype other than IgE which is induced by IL-4 (4). Indeed, the measurement of IgG4, which is produced in very small quantities, requires the utilization of IgG4 specific antibodies that are not employed in the RIA for IgG determination.

10

IL-12 induces the production of significant amounts of IFN- γ even in the presence of a high concentration of IL-4, that was shown to completely suppress IFN- γ production and to induce IgE synthesis in mixed lymphocyte cultures (6,7). Knowing that IFN- γ directs the
15 in vitro as well as the in vivo differentiation of naive T cells into TH1 type of cells, it is reasonable to assume that IL-12 may display the same activity even in the presence of IL-4, which may also be produced by non-T cells (26). According to this view, IL-12 might well pay a pivotal role in determining the outcome of certain
20 immune responses to certain antigens or pathogens that are known to preferentially generate TH1 or TH2 helper cells. The cellular origin of IL-12 is consistent with a putative role of this lymphokine in the differentiation of naive T cells. Indeed, IL-12 may be produced not only by Epstein-Barr virus transformed B cells from
25 which it was isolated but also by normal B cells that are known to be efficient antigen-presenting cells.

In preliminary experiments using neutralizing antibodies to IFN- γ the IL-12 mediated suppression of IgE synthesis by adult
30 PBMC was not consistently overcome. These results may be easily explained by (i) the relatively high levels of IFN- γ in IL-12 containing cultures and (ii) the difficulty in blocking the biological activity of endogenously produced IFN- γ . IL-12 also suppresses IgE by another mechanism which is IFN- γ independent. The existence of
35 such a mechanism is demonstrated by the ability of IL-12 to markedly inhibit IgE synthesis by IL-4 and hydrocortisone-

- costimulated neonatal lymphocytes which do not produce detectable amounts of IFN- γ . It is unlikely that such undetectable levels of IFN- γ (<1 IU) might nevertheless account for the suppression of IgE given that a very large excess of neutralizing anti-IFN- γ antibody failed to
- 5 increase the IgE response. Whereas both IL-12 and IFN- γ markedly suppress the accumulation of productive Ce mRNA in IL-4-stimulated PBMC (>90% suppression), IFN- γ , but not IL-12, also suppress the expression of germ-like transcript (50-70% inhibition). (5) IL-12 inhibits IgE synthesis by PBMC costimulated with IL-4 and anti-CD40
- 10 mAb, a model where IFN- γ was reported to be inactive (27). In three such experiments where PBMC were cultured with IL-4 and anti-CD40 mAb 89 (0.5 μ g/ml), the production of IgE dropped from 70 ± 28 ng/ml (mean \pm 1SD) to 20 ± 8 ng/ml in the presence of IL-12 (60pM) as compared to 79 ± 35 ng/ml in the presence of IFN- γ (100
- 15 IU/ml). Taken collectively, the present results demonstrate that, like the interferons, IL-12 plays an important role not only in protective immunity but also in the regulation of isotype selection.

TABLE 17

5	ADDITION	EXP. 1		EXP. 2	
		IgE	IFN- γ	IgE	IFN- γ
10	-	<0.2	214	<0.2	62
	IL-4	57	<1	204	<1
	IL-12	<0.2	3364	<0.2	2800
15	IL-4 + IL-12	19	1348	58	810

- 20 PBMC were cultured for 12 days in the absence or in the presence of IL-4 (10 ng/ml), IL-12 (60 pM) or both. Shown are the mean values of IgE (ng/ml) and IFN- γ (IU/ml) measured in 4 replicate cultures; the variation between the replicates was below 20%. Supernatant fluids from cultures of mock transfected COS cells were inactive when used at the same dilutions as the IL-12 containing supernatant
- 25 fluids (not shown).

TABLE 18
EFFECT OF IL-12 ON IgE SYNTHESIS BY NEONATAL LYMPHOCYTES STIMULATED WITH IL-4 AND
HYDROCORTISONE

ADDITION	EXP. 1		EXP. 2		EXP. 3		EXP. 4	
	IgE	IFN- γ	IgE	IFN- γ	IgE	IFN- γ	IgE	IFN- γ
	373	<1	44	<1	20	<1	302	<1
IL-12	6	<1	8	<1	<0.2	17	20	33
IL-12 + Anti-IFN- γ	7	<1	10	<1	<0.2	<1	21	<1
IL-12 + Anti-Lolp1	6	<1	NT	NT	NT	NT	18	128

Umbilical cord blood mononuclear cells were cultured for 12 days in the presence of IL-4 and 10 μ M hydrocortisone. IL-12 (60 pM), anti-IFN- γ mAb (1000 neutralizing U/ml) or the isotype-matched control (anti-Lolp1) mAb (50 μ g/ml) were added at the initiation of the culture. IgE (ng/ml) and IFN- γ (U/ml) were measured at day 12 and day 6 respectively. Shown are the mean values of quadruplicate cultures; the variation between the replicates was below 20%.

Example 16

Provided below are several exemplary formulations for parenteral use, injection, or aerosol administration.

5

The following formulations are parenteral solutions. These solutions may also be lyophilized to form powders.

3.0 mg/ml CLMF is dissolved in phosphate-buffered saline pH 7 (PBS pH 7) q.s. 1 ml. To this solution may be added one of the following: 0.2 mg/ml Polysorbate 80; or 5.0 mg human serum albumin; or 10.0 mg/ml benzyl alcohol. Alternatively, a formulation including 3.0 mg/ml CLMF; 25 mg/ml mannitol; and Tris buffer pH 7 q.s. 1 ml may be used.

15

A preferred formulation which is a lyophilized powder for injection includes 3.0 mg/ml CLMF, 3.0 mg/ml trehalose, and PBS pH 7 q.s. 1 ml.

20

All the above formulations are produced by mixing the indicated components in an appropriate vessel, filtering the resulting solution to sterility using an appropriate bacteria-retentive filter, and lyophilizing the resulting sterile solution in lyophilization vessels using an appropriate cycle. The lyophilization vessels should then be stoppered using the appropriate head pressure, gas, and stopper.

25

A suspension formulation for an inhalation aerosol may include the following components: 1.5 % w/w CLMF; 5.0 % w/w mannitol; 2.0 % w/w sorbitan trioleate; 64.0 % w/w Freon 12; 11.5 % w/w Freon 11; and 16.0 % w/w Freon 114. This formulation may be produced by conventional means. A preferred method of production is to prepare an aqueous solution of CLMF and mannitol and lyophilize this solution under suitable conditions. To the resulting dry product is added sorbitan trioleate and Freon 11, and the resulting mixture is

30

homogenized. The resulting suspension is filled into an aerosol container, which is crimped to seal with an appropriate valve inserted. The container is then pressure-filled with an 80:20 mixture of Freon 12 and Freon 114.

5

00101000 000000

REFERENCES

1. Finkelman, F.D., J. Holmes, I.M. Katona, J.F. Urban, Jr., J.P. Beckmann, L.S. Park, K.A. Schooley, R.L. Coffman, T.R. Mosmann, and W.E. Paul. 1990. Lymphokine control of in vivo immunoglobulin isotype selection. *Annu. Rev. Immunol.* 8:303.
2. Snapper, C.M., and W.E. Paul. 1987. Interferon- γ and B cell stimulatory factor-1 reciprocally regulate Ig isotype production. *Science (Washington, D.C.)* 236:944.
3. Pene, J., F. Rousset, F. Briere, I. Chretien, J.Y. Bonnefoy, H. Spits, T. Yokota, K. Arai, J. Banchereau, and J. de Vries. 1988. IgE production by normal human lymphocytes is induced by interleukin-4 and suppressed by interferons γ and α and prostaglandin E2. *Proc. Natl. Acad. Sci. USA* 85:6880.
4. Gascan, H., J.-F. Gauchat, M.-G. Roncarolo, H. Yssel, H. Spits, and J.E. deVries. 1991. Human B cell clones can be induced to proliferate and to switch to IgE and IgG4 synthesis by interleukin 4 and a signal provided by activated CD4⁺ T cell clones. *J. Exp. Med.* 173:747-750.
5. Gauchat, J.-F., D.A. Leberman, R.L. Coffman, H. Gascan, and J.E. deVries. 1990. Structure and expression of germlike ϵ transcripts in human B cells induced by interleukin 4 to switch to IgE production. *J. Exp. Med.* 172:463-473.
6. Peleman, R., J. Wu, C. Fargeas, and G. Delespesse. 1989. Recombinant interleukin 4 suppresses the production of interferon γ by human mononuclear cells. *J. Exp. Med.* 170:1751-1756.
7. Vercelli, D., H.H. Jabara, R. P. Lauener, and R.S. Geha. 1990. Interleukin-4 inhibits the synthesis of interferon- γ and

induces the synthesis of IgE in mixed lymphocyte cultures. J. Immunol. 144:570.

8. Coffman, R.L., K. Varkila, P. Scott, and R. Chatelain. 1991.
5 Role of cytokines in the differentiation of CD4⁺ T-cell subsets in vivo. Immunol. Reviews 123:189-207.
9. Gajewski, T.F., and F. W. Fitch. 1988. Anti-proliferative
effect of IFN- γ in immune regulation. I. IFN- γ inhibits the
10 proliferation of TH2 but not TH1 murine helper T cell clones. J. Immunol. 140:4245.
10. Fernandez-Botran, R., V.M. Sanders, T.R. Mosmann, J.W. Uhr, and E.S. Vilella. 1988. Lymphokine-mediated regulation of the
15 proliferative response of clones of TH1 and TH2 cells. J. Exp.Med. 168:543-548.
11. Scott, P. 1991. IFN- γ modulates the early development of
TH1 and TH2 responses in a murine model of cutaneous
20 leishmaniasis. J. Immunol. 147:3149-3155.
12. Finkelman, F.D., A. Svetic, I. Gresser, C. Snapper, J. Holmes, P.P. Trotta, I.M. Katona, and W.C. Gause. 1991. Regulation by
interferon- α of immunoglobulin isotype selection and lymphokine
25 production in mice. J.Exp.Med. 174:1179-1188.
13. Gubler, U., A.O. Chua, D.S. Schoenhaut, C.M. Dwyer, W. McComas, R. Motyka, N. Nabavi, A.G. Wolitzky, P.M. Quinn, P.C. Familletti, and M.K. Gately. 1991. Co-expression of two distinct
30 genes is required to generate secreted bioactive cytotoxic lymphocyte maturation factor. Proc. Natl. Acad. Sci. USA 88:4143.
14. Wolf, S.F., P.A. Temple, M. Kobayashi, D. Young, M. Dicig, L.Lowe, R. Dzialo, L. Fitz, C. Ferenz, R. M. Hewick, K. Kelleher, S.H. Herrmann, S.C. Clark, L. Azzoni, S.H. Chan, G. Trinchieri, and
35 B.Perussia. 1991. Cloning of cDNA for natural killer cell stimulatory

factor, a heterodimeric cytokine with multiple biologic effects on T and natural killer cells. *J. Immunol.* 146:3074-3081.

15. Kobayashi, M., L. Fitz, M. Ryan, R.M. Hewick, S.C. Clark, S. Chan, R. Loudon, F. Sherman, B. Perussia, and G. Trinchieri. 1989. Identification and purification of natural killer cell stimulatory factor (NKSF), a cytokine with multiple biologic effects on human lymphocytes. *J. Exp. Med.* 170:827-845.
- 10 16. Gately, M.K., B.B. Desai, A.G. Wolitzky, P.M. Quinn, C.M. Dwyer, F.J. Podlaski, P.C. Familletti, F. Sinigaglia, R. Chizzonite, U. Gubler, and A.S. Stern. 1991. Regulation of human lymphocyte proliferation by a heterodimeric cytokine, IL-12 (cytotoxic lymphocyte maturation factor). *J. Immunol.* 147:874-882.
- 15 17. Chan, S.H., B. Perussia, J.W. Gupta, M. Kobayashi, M. Pospisil, H.A. Young, S.F. Wolf, D. Young, S.C. Clark, and G. Trinchieri. 1991. Induction of interferon γ production by natural killer cell stimulatory factor: characterization of the responder cells and synergy with other inducers. *J. Exp. Med.* 173:869-879.
- 20 18. Rousset, F., E. Garcia, and J. Banchereau. 1991. Cytokine-induced proliferation and immunoglobulin production of human B lymphocytes triggered through their CD40 antigen. *J. Exp. Med.* 173:705-710.
- 25 19. Chizzonite, R., T. Truitt, F.J. Podlaski, A.G. Wolitzky, P.M. Quinn, P. Nunes, A.S. Stern, and M.K. Gately. 1991. IL-12: Monoclonal antibodies specific for the 40-kDa subunit block receptor binding and biologic activity on activated human lymphoblasts. *J. Immunol.* 147:1548-1556.
- 30 20. Sarfati, M., and G. Delespesse. 1988. Possible role of human lymphocyte receptor for IgE (CD23) or its soluble fragments in the in vitro synthesis of human IgE. *J. Immunol.* 141:2195.
- 35

21. Peleman, R., and G. Delespesse. 1990. In vitro synthesis of human IgE by neonatal lymphocytes: enhancing effect of interferon- γ . *Cell. Immunol.* 129:299-309.
- 5 22. Jabara, H.H., L.C. Schneider, S.K. Shapira, C. Alfieri, C.T. Moody, E. Kieff, R. Geha and D. Vercelli. 1990. Induction of germ-line and mature Ce transcripts in human B cells stimulated with rIL-4 and EBV. *J. Immunol.* 145:3468.
- 10 23. Kibler, R., M.J. Hicks, A.L. Wright, and L.M. Taussig. 1986. A comparative analysis of cord blood and adult lymphocytes: interleukin-2 and interferon production, natural killer cell activity, and lymphocyte populations. *Diagn.Immunol.* 4:201-208.
- 15 24. Wu, C.Y., M. Sarfati, C. Heusser, S. Fournier, M. Rubio-Trujillo, R. Peleman, and G. Delespesse. 1991. Glucocorticoids increase the synthesis of immunoglobulin E by interleukin 4-stimulated human lymphocytes. *J. Clin. Invest.* 87:870-877.
- 20 25. Vercelli, D. and R.S. Geha. 1991. Regulation of IgE synthesis in humans: A tale of two signals. *J. Aller. Clin. Immunol.* 88:285-295.
- 25 26. Plaut, M., J.H. Pierce, C.J. Watson, J. Hanley-Hyde, R.P. Nordan, W. E. Paul. 1989. Mast cell lines produce lymphokines in response to cross-linkage of Fc ϵ RI or calcium ionophores. *Nature* 339:64.
- 30 27. Zhang, K., E.A. Clark, and A. Saxon. 1991. CD40 stimulation provides an IFN- γ -independent and IL-4-dependent differentiation signal directly to human B cells for IgE production. *J. Immunol.* 146:1836-1842.

CLAIMS:

- 5 1. A process for producing Cytotoxic Lymphocyte
Maturation Factor (CMLF) in a substantially pure form comprising:
- stimulating lymphoblastoid cells to produce and secrete
cytokines into a supernatant liquid;
- 10 collecting the supernatant liquid produced by the
stimulated cells;
- separating the supernatant liquid into protein fractions;
- 15 testing each protein fraction for the presence of CLMF by
means of a suitable assay;
- retaining the protein fractions which contain CLMF;
- 20 isolating the CLMF from said CLMF-containing protein
fraction into a substantially pure form.
2. The process of Claim 1 wherein the cells are NC-37 B
- 25 lymphoblastoid cells.
3. The process of Claim 1 wherein the supernatant liquid is
separated into protein fractions by means of a strong cation exchange
column.
- 30 4. The process of Claim 3 wherein the strong cation
exchange is a sulfopropyl cation exchange.
5. The process of Claim 3, further comprising passing the
- 35 protein fractions containing CLMF through an agarose gel, said
agarose gel being comprised of a Blue B dye covalently coupled to an

agarose matrix; and testing the protein fractions which are eluted through said agarose gel for the presence of CLMF and retaining the protein fractions which contain CLMF.

5 6. The process of Claim 5 further comprising eluting the protein fractions containing CLMF of Claim 7 through a strong anion exchange in a High Performance Liquid Chromatography or a Fast Protein Liquid Chromatography mode to obtain protein fractions;

10 testing the protein fractions eluted through the strong anion exchange for the presence of CLMF and retaining the protein fraction containing CLMF.

15 7. A protein comprising Cytotoxic Lymphocyte Maturation Factor (CLMF) in a substantially pure form or a protein which exhibits CLMF activity and contains a biologically active portion of the amino acid sequence of CLMF or which contains an amino acid sequence of CLMF as well as other amino acids or proteins containing analogous sequences to CLMF or its biologically active fragments
20 which proteins exhibit CLMF activity.

8. The protein of Claim 7, wherein CLMF is comprised of a 75 kDa polypeptide, or biologically active fragment thereof.

9. The protein of Claim 8, wherein the 75 kDa polypeptide has an amino acid composition comprised of:

	<u>Amino acid</u>	<u>mol %</u>
5	Aspartic acid or asparagine	10.8
	Threonine	7.2
	Serine	8.9
10	Glutamic acid or glutamine	13.1
	Proline	3.8
	Glycine	4.7
	Alanine	5.9
	Cysteine	2.9
15	Valine	6.2
	Methionine	1.9
	Isoleucine	4.2
	Leucine	9.4
	Tyrosine	3.6
20	Phenylalanine	3.7
	Histidine	1.8
	Lysine	7.7
	Arginine	4.4
25	Tryptophan	ND

10. The protein of Claim 9, wherein the 75 kDa polypeptide is comprised of two disulfide linked subunits, a 35 kDa subunit and a 40 kDa subunit.

30

11. The protein of Claim 10, wherein the 35 kDa subunit polypeptide has the following amino acid sequence.

Arg Asn Leu Pro Val Ala Thr Pro Asp Pro Gly MET Phe Pro Cys Leu
5 His His Ser Gln Asn Leu Leu Arg Ala Val Ser Asn MET Leu Gln Lys
Ala Arg Gln Thr Leu Glu Phe Tyr Pro Cys Thr Ser Glu Glu Ile Asp
His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val Glu Ala Cys Leu
Pro Leu Glu Leu Thr Lys Asn Glu Ser Cys Leu Asn Ser Arg Glu Thr
Ser Phe Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe
10 MET MET Ala Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys MET Tyr
Gln Val Glu Phe Lys Thr MET Asn Ala Lys Leu Leu MET Asp Pro Lys
Arg Gln Ile Phe Leu Asp Gln Asn MET Leu Ala Val Ile Asp Glu Leu
MET Gln Ala Leu Asn Phe Asn Ser Glu Thr Val Pro Gln Lys Ser Ser
Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys Ile Lys Leu Cys Ile Leu
15 Leu His Ala Phe Arg Ile Arg Ala Val Thr Ile Asp Arg Val Thr Ser
Tyr Leu Asn Ala Ser

12. The protein of Claim 10, wherein the 40 kDa subunit polypeptide has the following amino acid sequence:

20 Ile Trp Glu Leu Lys Lys Asp Val Tyr Val Val Glu Leu Asp Trp Tyr Pro Asp
Ala Pro Gly Glu MET Val Val Leu Thr Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp
Thr Leu Asp Gln Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
25 Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val Leu Ser His Ser
Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp Ile Leu Lys Asp Gln
Lys Glu Pro Lys Asn Lys Thr Phe Leu Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe
Thr Cys Trp Trp Leu Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser Arg
Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser Ala Glu Arg Val
Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu Cys Gln Glu Asp Ser Ala Cys Pro
30 Ala Ala Glu Glu Ser Leu Pro Ile Glu Val MET Val Asp Ala Val His Lys Leu Lys Tyr
Glu Asn Tyr Thr Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn
Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp Glu Tyr Pro Asp
Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr Phe Cys Val Gln Val Gln Gly Lys
Ser Lys Arg Glu Lys Lys Asp Arg Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys
35 Arg Lys Asn Ala Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser
Glu Trp Ala Ser Val Pro Cys Ser

13. The protein of Claim 8 wherein said protein is a biologically active fragment.

40

14. An isolated polynucleotide encoding for CLMF or a biologically active fragment thereof.

15. A recombinant vector containing the polynucleotide of Claim 14.

16. A host cell transformed with the vector of Claim 15.

17. An isolated polynucleotide encoding for a 40 kDa subunit of CLMF having the following DNA sequence:

ATG TGT CAC CAG CAG TTG GTC ATC TCT TGG TTT TOC CTG GTT TTT CTG GCA TCT CCC CTC
15 GTG GCC ATA TGG GAA CTG AAG AAA GAT GTT TAT GTC GTA GAA TTG GAT TGG TAT CCG GAT
GCC CCT GGA GAA ATG GTG GTC CTC ACC TGT GAC ACC CCT GAA GAA GAT GGT ATC ACC TGG
ACC TTG GAC CAG AGC AGT GAG GTC TTA GGC TCT GGC AAA ACC CTG ACC ATC CAA GTC AAA
GAG TTT GGA GAT GCT GGC CAG TAC ACC TGT CAC AAA GGA GGC GAG GTT CTA AGC CAT TCG
CTC CTG CTG CTT CAC AAA AAG GAA GAT GGA ATT TGG TCC ACT GAT ATT TTA AAG GAC CAG
20 AAA GAA CCC AAA AAT AAG ACC TTT CTA AGA TGC GAG GCC AAG AAT TAT TCT GGA CGT TTC
ACC TGC TGG TGG CTG ACG ACA ATC AGT ACT GAT TTG ACA TTC AGT GTC AAA AGC AGC AGA
GGC TCT TCT GAC CCC CAA GGG GTG ACG TGC GGA GCT GCT ACA CTC TCT GCA GAG AGA GTC
AGA GGG GAC AAC AAG GAG TAT GAG TAC TCA GTG GAG TGC CAG GAG GAC AGT GCC TGC CCA
GCT GCT GAG GAG AGT CTG CCC ATT GAG GTC ATG GTG GAT GCC GTT CAC AAG CTC AAG TAT
25 GAA AAC TAC ACC AGC AGC TTC TTC ATC AGG GAC ATC ATC AAA CCT GAC CCA CCC AAG AAC
TTG CAG CTG AAG CCA TTA AAG AAT TCT CCG CAG GTG GAG GTC AGC TGG GAG TAC CCT GAC
ACC TGG AGT ACT CCA CAT TCC TAC TTC TOC CTG ACA TTC TGC GTT CAG GTC CAG GGC AAG
AGC AAG AGA GAA AAG AAA GAT AGA GTC TTC ACG GAC AAG ACC TCA GCC ACG GTC ATC TGC
CGC AAA AAT GCC AGC ATT AGC GTG CCG GGC CAG GAC CGC TAC TAT AGC TCA TCT TGG AGC
30 GAA TGG GCA TCT GTG CCC TGC AGT

18. A recombinant vector containing the polynucleotide of claim 17 or fragment thereof.

19. A host cell transformed with the vector of claim 18.

20. An isolated polynucleotide encoding for a 35 kDa subunit polypeptide of CLMF having the following DNA sequence:

5 ATG TGT CCA GCG CGC AGC CTC CTC CTT GTG GCT ACC CTG GTC CTC CTG
GAC CAC CTC AGT TTG GCC AGA AAC CTC CCC GTG GGC ACT CCA GAC CCA
GGA ATG TTC CCA TGC CTT CAC CAC TCC CAA AAC CTG CTG AGG GGC GTC
AGC AAC ATG CTC CAG AAG GCC AGA CAA ACT CTA GAA TTT TAC CCT TGC
10 ACT TCT GAA GAG ATT GAT CAT GAA GAT ATC ACA AAA GAT AAA ACC AGC
ACA GTG GAG GCC TGT TTA CCA TTG GAA TTA ACC AAG AAT GAG AGT TGC
CTA AAT TCC AGA GAG ACC TCT TTC ATA ACT AAT GGG AGT TGC CTG GGC
TCC AGA AAG ACC TCT TTT ATG ATG GCC CTG TGC CTT AGT AGT ATT TAT
GAA GAC TTG AAG ATG TAC CAG GTG GAG TTC AAG ACC ATG AAT GCA AAG
CTT CTG ATG GAT CCT AAG AGG CAG ATC TTT CTA GAT CAA AAC ATG CTG
15 GCA GTT ATT GAT GAG CTG ATG CAG GCC CTG AAT TTC AAC AGT GAG ACT
GTG CCA CAA AAA TCC TCC CTT GAA GAA CCG GAT TTT TAT AAA ACT AAA
ATC AAG CTC TGC ATA CTT CTT CAT GCT TTC AGA ATT CCG GCA GTG ACT
ATT GAC AGA GTG ACG AGC TAT CTG AAT GCT TCC

20 21. A recombinant vector containing the isolated polynucleotide of claim 20.

22. A host cell transformed with the vector of claim 21.

25 23. A method for stimulating LAK cells comprising:

30 treating the LAK cells with CLMF or a biologically active fragment of CLMF and with IL2 or a biologically active fragment thereof.

24. A method for stimulating activated T-cells comprising:
35 treating T-cells with CLMF or a biologically active fragment of CLMF.

25. A method for stimulating activated T-cells as recital in Claim 24 further comprising:

40

treating the activated T-cells with IL2 or a biologically active fragment thereof.

26. A method for stimulating Natural Killer Cells comprising:

5

treating the Natural Killer Cells with CLMF or a biologically active fragment of CLMF.

27. Antibodies in a substantially pure form to CLMF.

10

28. Antibodies to CLMF of Claim 27 wherein the antibodies are polyclonal.

15

29. Antibodies to CLMF of Claim 27 wherein the antibodies are monoclonal.

30. A process for producing Cytotoxic Lymphocyte Maturation Factor (CMLF) in a substantially pure form comprising:

20

stimulating cells capable of producing CLMF to produce and secrete cytokines into a supernatant liquid;

25

collecting the supernatant liquid produced by the stimulated cells;

separating the supernatant liquid into protein fractions;

30

testing each protein fraction for the presence of CLMF by means of a suitable assay;

retaining the protein fractions which contain CLMF;

35

isolating the CLMF from said CLMF-containing protein fraction into a substantially pure form.

31. A process of producing substantially pure CLMF from supernatant liquids obtained from culturing cells, which liquid contains CLMF together with other proteins comprising:

- 5 separating the supernatant liquid into protein fractions;

 testing each protein fraction for the presence of CLMF by means of a suitable assay;
- 10 retaining the protein fractions which contain CLMF;

 isolating the CLMF from said CLMF-containing protein fraction into a substantially pure form.

15 32. A process for producing substantially pure CLMF for liquids containing CLMF together with other proteins comprising:

- separating the supernatant liquid into protein fractions;
- 20 testing each protein fraction for the presence of CLMF by means of a suitable assay;

 retaining the protein fractions which contain CLMF;
- 25 isolating the CLMF from said CLMF-containing protein fraction into a substantially pure form.


ABSTRACT OF THE DISCLOSURE


The present invention is a novel cytokine protein called IL-12 or Cytotoxic Lymphocyte Maturation Factor (CLMF) which is
5 produced and synthesized by human NC-37 B lymphoblastoid cells (American Type Culture Collection, Rockville, MD). CLMF synergistically induces with low concentrations of IL-2 the cytolytic activity of Lymphokine Activated Killer (LAK) cells, and CLMF is capable of stimulating T-cell growth.

10

Also claimed are the cloned gene for CLMF, its recombination in a suitable vector, the transformed cells containing said vector, the recombinant protein produced by the transformed cells and antibodies to CLMF.

15

**FIG. 1**

**FIG. 2**





FIG. 3

**FIG. 4**




FIG. 5

00401000, 000000





FIG. 6

**FIG. 7**





FIG. 8






FIG. 10




FIG. 11






FIG. 12

**FIG. 13**

**FIG. 14**




FIG. 15





FIG. 16

10 20 30 40
 I-W-E-L-K-D-V-V-V-E-L-D-W-Y-P-D-A-P-G-E-M-V-V-L-T-C-D-T-P-E-E-D-G-I-T-W-T-L
 N-t t-----f60-----t
 Tr t-----f60-----t
 V8 t-t47-t-----f54-----t-----f57-----t
 50 60 70
 D-Q-S-S-E-V-L-G-S-G-K-I-L-T-I-Q-V-K-E-F-G-D-A-G-Q-Y-T-C-H-K-G-G-E-V-L-S-H-S-L-L
 Tr t-----f57-----t
 V8 t---f57-----t-----f78-----t
 90 100 110 120
 L-L-H-K-K-E-D-G-I-W-S-T-D-I-L-K-D-Q-K-E-P-K-N-K-T-F-L-R-C-E-A-K-N-Y-S-G-R-F-T-C
 Tr t-----f67-----t-----f52-----t
 Tr t-----f67-----t-----f52-----t
 130 140 150 160
 W-W-L-T-T-I-S-I-D-I-F-S-V-K-S-S-R-Q-S-D-P-Q-G-V-T-C-Q-A-A-T-L-S-A-E-R-V-R-G
 Tr t-----f55-----t
 170 180 190 200
 D-N-K-E-Y-E-Y-S-V-E-C-Q-E-D-S-A-C-P-A-A-E-E-S-L-P-I-E-V-M-V-D-A-V-H-K-L-K-Y-E-N
 Tr t-----f70-----t
 210 220 230 240
 Y-I-S-S-F-I-R-D-I-I-K-P-D-P-K-N-L-Q-I-K-P-L-K-N-S-R-Q-V-E-V-S-W-E-Y-P-D-T-W
 Tr t-----f70-----t
 250 260 270 280
 S-T-P-H-S-Y-F-S-L-T-F-C-V-Q-G-K-S-K-R-E-K-K-D-R-V-F-I-D-K-T-S-A-T-V-I-C-R-K
 Tr t-----f67-----t-----f55&f57-----t
 290 300 306
 N-A-S-I-S-V-R-A-Q-D-R-Y-Y-S-S-S-W-S-E-W-A-S-V-P-C-S
 Tr t-----f67-----t
 Tr t-----f72-----t-----f72-----t

FIG. 17

**FIG. 18**




FIG. 19




FIG. 20




FIG. 21





FIG. 22




FIG. 23




FIG. 24

10			20			30			40			49			58					
GTTTCAGGGC CATTTGGACTC TCCGTCCTGC CCAGAGCAAG									>											
ATG			TGT			CAC			CAG			CAG			TTG			GTC		
MET			Cys			His			Gln			Gln			Leu			Val		
67			76			85			94			103			112					
ATC	TCT	TGG	TTT	TCC	CTG	GTT	TTT	CTG	GCA	TCT	CCC	CTC	GTG	GCC	ATA	TGG	GAA			
Ile	Ser	Trp	Phe	Ser	Leu	Val	Phe	Leu	Ala	Ser	Pro	Leu	Val	Ala	Ile	Trp	Glu			
121			130			139			148			157			166					
CTG	AAG	AAA	GAT	GTT	TAT	GTC	GTA	GAA	TTG	GAT	TGG	TAT	CCG	GAT	GCC	CCT	GGA			
Leu	Lys	Lys	Asp	Val	Tyr	Val	Val	Glu	Leu	Asp	Trp	Tyr	Pro	Asp	Ala	Pro	Gly			
175			184			193			202			211			220					
GAA	ATG	GTG	GTC	CTC	ACC	TGT	GAC	ACC	CCT	GAA	GAA	GAT	GGT	ATC	ACC	TGG	ACC			
Glu	MET	Val	Val	Leu	Thr	Cys	Asp	Thr	Pro	Glu	Glu	Asp	Gly	Ile	Thr	Trp	Thr			
229			238			247			256			265			274					
TTG	GAC	CAG	AGC	AGT	GAG	GTC	TTA	GGC	TCT	GGC	AAA	ACC	CTG	ACC	ATC	CAA	GTC			
Leu	Asp	Gln	Ser	Ser	Glu	Val	Leu	Gly	Ser	Gly	Lys	Thr	Leu	Thr	Ile	Gln	Val			
283			292			301			310			319			328					
AAA	GAG	TTT	GGA	GAT	GCT	GGC	CAG	TAC	ACC	TGT	CAC	AAA	GGA	GGC	GAG	GTT	CTA			
Lys	Glu	Phe	Gly	Asp	Ala	Gly	Gln	Tyr	Thr	Cys	His	Lys	Gly	Gly	Glu	Val	Leu			

FIG. 25a

337	346	355	364	373	382
<u>AGC</u> <u>CAT</u> <u>TCG</u> <u>CTC</u> <u>CTG</u> <u>CTT</u> <u>CAC</u> <u>AAA</u> <u>AAG</u> <u>GAA</u> <u>GAT</u> <u>GGA</u> <u>ATT</u> <u>TGG</u> <u>TCC</u> <u>ACT</u> <u>GAT</u>					
Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp Ser Thr Asp					
391	400	409	418	427	436
<u>ATT</u> <u>TTA</u> <u>AAG</u> <u>GAC</u> <u>CAG</u> <u>AAA</u> <u>GAA</u> <u>CCC</u> <u>AAA</u> <u>AAT</u> <u>AAG</u> <u>ACC</u> <u>TTT</u> <u>CTA</u> <u>AGA</u> <u>TGC</u> <u>GAG</u> <u>GCC</u>					
Ile Leu Lys Asp Gln Lys Lys Glu Pro Lys Asn Lys Thr Phe Leu Arg Cys Glu Ala					
445	454	463	472	481	490
<u>AAG</u> <u>AAT</u> <u>TAT</u> <u>TCT</u> <u>GGA</u> <u>CGT</u> <u>TTC</u> <u>ACC</u> <u>TGC</u> <u>TGG</u> <u>CTG</u> <u>ACG</u> <u>ACA</u> <u>ATC</u> <u>AGT</u> <u>ACT</u> <u>GAT</u>					
Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp Leu Thr Thr Ile Ser Thr Asp					
499	508	517	526	535	544
<u>TTG</u> <u>ACA</u> <u>TTT</u> <u>AGT</u> <u>GTC</u> <u>AAA</u> <u>AGC</u> <u>AGC</u> <u>AGA</u> <u>GGC</u> <u>TCT</u> <u>TCT</u> <u>GAC</u> <u>CCC</u> <u>CAA</u> <u>GGG</u> <u>GTG</u> <u>ACG</u>					
Leu Thr Phe Ser Val Lys Ser Ser Arg Gly Ser Ser Asp Pro Gln Gly Val Thr					
553	562	571	580	589	598
<u>TGC</u> <u>GGA</u> <u>GCT</u> <u>GCT</u> <u>ACA</u> <u>CTC</u> <u>TCT</u> <u>GCA</u> <u>GAG</u> <u>AGA</u> <u>GTC</u> <u>AGA</u> <u>GGG</u> <u>GAC</u> <u>ARC</u> <u>ARG</u> <u>GAG</u> <u>TAT</u>					
Cys Gly Ala Ala Thr Leu Ser Ala Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr					
607	616	625	634	643	652
<u>GAG</u> <u>TAC</u> <u>TCA</u> <u>CTG</u> <u>GAG</u> <u>TGC</u> <u>CAG</u> <u>GAG</u> <u>GAC</u> <u>AGT</u> <u>GCC</u> <u>TGC</u> <u>CCA</u> <u>GCT</u> <u>GCT</u> <u>GAG</u> <u>GAG</u> <u>AGT</u>					
Glu Tyr Ser Val Glu Cys Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu Ser					

FIG. 25b

661	670	679	688	697	706												
<u>CTG</u> <u>CCC</u> <u>ATT</u> <u>GAG</u> <u>GTC</u> <u>ATG</u> <u>GAT</u> <u>GCC</u> <u>GTT</u> <u>CAC</u> <u>AAG</u> <u>CTC</u> <u>AAG</u> <u>TAT</u> <u>GAA</u> <u>AAC</u> <u>TAC</u>																	
Leu	Pro	Ile	Glu	Val	Met	Val	His	Lys	Leu	Lys	Tyr	Glu	Asn	Tyr			
715	724	733	742	751	760												
<u>ACC</u> <u>AGC</u> <u>AGC</u> <u>ATC</u> <u>TTC</u> <u>ATC</u> <u>AGG</u> <u>GAC</u> <u>ATC</u> <u>ATC</u> <u>AAA</u> <u>CCT</u> <u>GAC</u> <u>CCA</u> <u>CCC</u> <u>AAG</u> <u>AAC</u> <u>TTG</u>																	
Thr	Ser	Ser	Phe	Phe	Ile	Arg	Asp	Ile	Ile	Lys	Pro	Asp	Pro	Pro	Lys	Asn	Leu
769	778	787	796	805	814												
<u>CAG</u> <u>CTG</u> <u>AAG</u> <u>CCA</u> <u>TTA</u> <u>AAG</u> <u>AAT</u> <u>TCT</u> <u>CGG</u> <u>CAG</u> <u>GTC</u> <u>GAG</u> <u>GTC</u> <u>AGC</u> <u>TGG</u> <u>GAG</u> <u>TAC</u> <u>CCT</u>																	
Gln	Leu	Lys	Pro	Leu	Lys	Asn	Ser	Arg	Gln	Val	Glu	Val	Ser	Trp	Glu	Tyr	Pro
823	832	841	850	859	868												
<u>GAC</u> <u>ACC</u> <u>TGG</u> <u>AGT</u> <u>ACT</u> <u>CCA</u> <u>CAT</u> <u>TCC</u> <u>TAC</u> <u>TTT</u> <u>TCC</u> <u>CTG</u> <u>ACA</u> <u>TTC</u> <u>TGC</u> <u>GTT</u> <u>CAG</u> <u>GTC</u>																	
Asp	Thr	Trp	Ser	Thr	Pro	His	Ser	Tyr	Phe	Ser	Leu	Thr	Phe	Cys	Val	Gln	Val
877	886	895	904	913	922												
<u>CAG</u> <u>GGC</u> <u>AAG</u> <u>AGC</u> <u>AAG</u> <u>AGA</u> <u>GAA</u> <u>AAA</u> <u>GAT</u> <u>AGA</u> <u>GTC</u> <u>TTT</u> <u>ACG</u> <u>GAC</u> <u>AAG</u> <u>ACC</u> <u>TCA</u>																	
Gln	Gly	Lys	Ser	Lys	Arg	Glu	Lys	Lys	Asp	Arg	Val	Phe	Thr	Asp	Lys	Thr	Ser
931	940	949	958	967	976												
<u>GGC</u> <u>ACG</u> <u>GTC</u> <u>ATC</u> <u>TGC</u> <u>CGC</u> <u>AAA</u> <u>AAT</u> <u>GCC</u> <u>AGC</u> <u>ATT</u> <u>AGC</u> <u>GTG</u> <u>CGG</u> <u>CCC</u> <u>CAG</u> <u>GAC</u> <u>CGC</u>																	
Ala	Thr	Val	Ile	Cys	Arg	Lys	Asn	Ala	Ser	Ile	Ser	Val	Arg	Ala	Gln	Asp	Arg

FIG. 25c

985 994 1003 1012 1021 1034
TAC TAT AGC TCA TCT TGG AGC GAA TGG GCA TCT GTG CCC TGC AGT TAGTTCTCA
 Tyr Tyr Ser Ser Trp Ser Glu Trp Ala Ser Val Pro Cys Ser
 1044 1054 1064 1074 1084 1104
 TCCAGGATGA AATTTCGAG GAAAGTGA AGATATTAG CAAATGTTT AAAGACCAA CGGATAGAC
 1114 1124 1134 1144 1154 1174
 CCAAAAGAT AATTCTATC TCATTGCTT TAAAGCTT TTTAGGATC ACAATGATAT CTTTGCTGA
 1184 1194 1204 1214 1224 1244
 TTGTATAGT TAGATGCTTA ATGCTCATG AAACAATCG CTATTTCG TATAGATTT CCAGCTCTCA
 1254 1264 1274 1284 1294 1314
 AGTTGCCATG GGCCTTCATG CTATTAAAT ATTAAAGTAA TTTATGTAAT TATTAGTATA TTACTGTAT
 1324 1334 1344 1354 1364 1374 1384 1394
 TTAAGTTTG TCTGCCAGGA TGTATGGAAT GTTTCATCT CTATAGCCT GATCATCAG GATCAGTCCC TATTATGCA AAT

FIG. 25d

10	20	30	40	50	60	70
GAATTCACAG AAAGCAAGAG ACCAGAGTCC CGGGAAAGTC CTGCCGCGCC TCGGGACAAT TATAAAAATG						
80	90	100	110	120	130	140
TGGCCCCCTG GGTGAGCCTC CCAGCCACCG CCTTCACCTG CCGCGGGCAC AGGTCTGCAT CCAGCGGCTC						
150	160	169	178	187	196	
GCCCTGTGTC CCTGCAGTGC CGGCTCAGC ^{>} ATG TGT CCA GCG CGC AGC CTC CTC CTT GTG MET Cys Pro Ala Arg Ser Leu Leu Leu Val						
205	214	223	232	241	250	
GCT ACC CTG GTC CTC CTG GAC CAC CTC AGT TTG GCC AGA AAC CTC CCC GTG GCC Ala Thr Leu Val Leu Leu Asp His Leu Ser Leu Ala Arg Asn Leu Pro Val Ala						
259	268	277	286	295	304	
ACT CCA GAC CCA GGA ATG TTC CCA TGC CTT CAC CAC TCC CAA AAC CTG CTG AGG Thr Pro Asp Pro Gly MET Phe Pro Cys Leu His His Ser Gln Asn Leu Leu Arg						
313	322	331	340	349	358	
GCC GTC AGC AAC ATG CTC CAG AAG GCC AGA CAA ACT CTA GAA TTT TAC CCT TGC Ala Val Ser Asn MET Leu Gln Lys Ala Arg Gln Thr Leu Glu Phe Tyr Pro Cys						
367	376	385	394	403	412	
ACT TCT GAA GAG ATT GAT CAT GAA GAT ATC ACA AAA GAT AAA ACC AGC ACA GTG Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp Lys Thr Ser Thr Val						

FIG. 26a

421	430	439	448	457	466
GAG GCC TGT TTA CCA TTG GAA TTA ACC AAG AAT GAG AGT TGC CTA AAT TCC AGA	Glu Ala Cys Leu Pro Leu Glu Leu Thr Lys Asn Glu Ser Cys Leu Asn Ser Ser Arg				
475	484	493	502	511	520
GAG ACC TCT TTC ATA ACT AAT GGG AGT TGC CTG GCC TCC AGA AAG ACC TCT TTT	Glu Thr Ser Phe Ile Thr Asn Gly Ser Cys Leu Ala Ser Arg Lys Thr Ser Phe				
529	538	547	556	565	574
ATG ATG GCC CTG TGC CTT AGT AGT ATT TAT GAA GAC TTG AAG ATG TAC CAG GTG	MET MET Ala Leu Cys Leu Ser Ser Ile Tyr Glu Asp Leu Lys MET Tyr Gln Val				
583	592	601	610	619	628
GAG TTC AAG ACC ATG AAT GCA AAG CTT CTG ATG GAT CCT AAG AGG CAG ATC TTT	Glu Phe Lys Thr MET Asn Ala Lys Leu Leu MET Asp Pro Lys Arg Gln Ile Phe				
637	646	655	664	673	682
CTA GAT CAA AAC ATG CTG GCA GTT ATT GAT GAG CTG ATG CAG GCC CTG AAT TTC	Leu Asp Gln Asn MET Leu Ala Val Ile Asp Glu Leu MET Gln Ala Leu Asn Phe				
691	700	709	718	727	736
AAC AGT GAG ACT GTG CCA CAA AAA TCC TCC CTT GAA GAA CCG GAT TTT TAT AAA	Asn Ser Glu Thr Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe Tyr Lys				
745	754	763	772	781	790
ACT AAA ATC AAG CTC TGC ATA CTT CTT CAT GCT TTC AGA ATT CGG GCA GTG ACT	Thr Lys Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg Ala Val Thr				

FIG. 26b

799		808		817		826				
<u>ATT</u>	<u>GAC</u>	<u>AGA</u>	<u>GTG</u>	<u>ACG</u>	<u>AGC</u>	<u>TAT</u>	<u>CTG</u>	<u>AAT</u>	<u>GCT</u>	<u>TCC</u>
Ile	Asp	Arg	Val	Thr	Ser	Tyr	Leu	Asn	Ala	Ser

836	846	856
TAAAAAGCGA GGTCCCTCCA AACCGTTGTC		

FIG. 26c




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35





FIG. 36

**FIG. 37**





FIG. 38

**FIG. 39**

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: GATELY, MAURICE K. et al.

Serial No.: To Be Assigned
(Division Of Serial No.
08/459,151)

Group Art Unit: To Be
Assigned

Examiner: To Be Assigned

Filed: On Even Date Herewith

Attorney Docket No.: 1803-247

For: PURIFICATION AND
CHARACTERIZATION OF
CYTOTOXIC LYMPHOCYTE
MATURATION FACTOR AND
MONOCLONAL ANTIBODIES
THERETO

DECLARATION OF THOMAS E. FRIEBEL
UNDER 37 C.F.R. § 1.608(a)

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

I, Thomas E. Friebe, declare and state as follows:

1. I am a partner at the law firm of Pennie & Edmonds LLP and am an attorney of record in the above-captioned application.
2. I am familiar with the proposed count in the accompanying "Request Under 37 C.F.R. §§ 1.607 And 1.608(a) For Interference With A Patent."

3. There is a basis upon which Applicants are entitled to a judgment relative to the patentee of United States Patent No. 5,811,523.

I declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date September 22, 1999



Thomas E. Friebe

Declaration and Power of Attorney For Patent Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

PURIFICATION AND CHARACTERIZATION OF CYTOTOXIC LYMPHOCYTE MATURATION FACTOR
AND MONOCLONAL ANTIBODIES THERETO

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and was amended on _____
(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

(Number)

(Country)

(Day/Month/Year Filed)

☐ Yes ☐ No

(Number)

(Country)

(Day/Month/Year Filed)

☐ Yes ☐ No

(Number)

(Country)

(Day/Month/Year Filed)

☐ Yes ☐ No

45 91 51

I hereby claim the benefit under 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

07/455,708	December 22, 1989	Abandoned
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)
07/520,935	May 9, 1990	Pending
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)
07/572,284	August 27, 1990	Pending
(Application Serial No.)	(Filing Date)	(Status)
		(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

George M. Gould	(Reg. No. 20970)	William H. Epstein	(Reg. No. 20008)
Bernard S. Leon	(Reg. No. 20756)	William G. Isgro	(Reg. No. 22041)
		George W. Johnston	(Reg. No. 28090)
		Catherine R. Roseman	(Reg. No. 34240)

Send Correspondence to:

George M. Gould, Esq., Hoffmann-La Roche Inc., 340 Kingsland Street, Nutley, New Jersey 07110-1199

Direct Telephone Calls to: (name and telephone number)

Catherine R. Roseman at telephone No. (201) 235-6208

Maurice Kent Gately

Full name of sole or first inventor

Maurice Kent Gately

Inventor's signature

Date

Montville, Morris County, New Jersey

Residence

USA

Citizenship

162 Konner Avenue, Montville, NJ 07058

Post Office Address

Ulrich Andreas Gubler

Full name of second joint inventor, if any

Ulrich Andreas Gubler

Second inventor's signature

Date

Glen Ridge, Essex County, New Jersey

Residence

Switzerland

Citizenship

4 Imness Place, Glen Ridge, NJ 07028

Post Office Address

(Supply similar information and signature for third and subsequent joint inventors.)

45 91 51

Jeffrey David Hulmes

Full name of third joint inventor, if any

Jeffrey David Hulmes

3/22/92
Date

Third Inventor's signature

Ringwood, Passaic County, New Jersey

Residence

USA

Citizenship

23 Welch Road, Ringwood, NJ 07456

Post Office Address

Frank John Podlaski

Full name of fourth joint inventor, if any

Frank John Podlaski

3-18-92
Date

Fourth Inventor's signature

New City, Rockland County, New York

Residence

USA

Citizenship

28 Lombardi Drive, New City, NY 10956

Post Office Address

Alvin Seth Stern

Alvin Seth Stern

Full name of fifth joint inventor, if any

Fifth Inventor's signature

Passaic Park, Passaic County, New Jersey

Date

Residence

USA

Citizenship

295 Brook Avenue, Passaic Park, NJ 07055

Post Office Address

Richard Anthony Chi

Full name of sixth joint inventor, or, if any

Richard Anthony Chi
6th Inventor's signature
South Kent, Litchfield County, Connecticut

3-18-92

Date

Residence

USA

Citizenship

104 Richards Road, South Kent, Connecticut 06785

Post Office Address

Yu-Ching Eugene Pan

Full name of seventh joint inventor, if any

Yu-Ching Eugene Pan
7th Inventor's signature
Pine Brook, Essex County, New Jersey

3/18/92

Date

Residence

USA

Citizenship

10 Crane Drive, Pine Brook, NJ 07058

Post Office Address

Title 37, Code of Federal Regulations, §1.56

Duty to disclose information material to patentability.

- (a) Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is cancelled or withdrawn from consideration, or the application becomes abandoned.
- (b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and
- (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim: or
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
- (i) Opposing an argument of unpatentability relied on by the Office, or
- (ii) Asserting an argument of patentability.

EXPRESS MAIL NO.: EM061 021 888US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: Gately et al.

Application No.: 08/459,151

Art Unit: 1646

Filed: June 2, 1995

Examiner: P. Mertz, Ph.D.

For: PURIFICATION AND
CHARACTERIZATION OF
CYTOTOXIC LYMPHOCYTE
MATURATION FACTOR AND
MONOCLONAL ANTIBODIES
THERE TO

Attorney Docket No.: 1803-241

ASSOCIATE POWER OF ATTORNEYAssistant Commissioner for Patents
Washington, D.C. 20231


Sir:

Please recognize Thomas E. Friebe (Reg. No. 29,258) as Associate Attorney and
Nikolaos C. George (Reg. No. 39,201) as Associate Agent in this application and any
divisional and/or any continuations thereof.

Kindly continue to address all communications to the attorney of record:

George W. Johnston
340 Kingsland Street
Nutley, New Jersey 07110

Respectfully submitted,

Date September 22, 1999
20,008
William H. Epstein (Reg. No.)
Attorney for Applicant(s)
Telephone: (973) 235-3723
Telefax: (973) 235-2363

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: GATELY, MAURICE K. et al.

Serial No.: To Be Assigned
(Division Of Serial No.
08/459,151)Group Art Unit: To Be
Assigned

Examiner: To Be Assigned

Filed: On Even Date Herewith

Attorney Docket No.: 1803-247

For: PURIFICATION AND
CHARACTERIZATION OF
CYTOTOXIC LYMPHOCYTE
MATURATION FACTOR AND
MONOCLONAL ANTIBODIES
THERETODECLARATION OF DR. DAVID H. PRESKYAssistant Commissioner for Patents
Washington, D.C. 20231

Sir:

I, DAVID H. PRESKY, PH.D., do declare that:

1. I am currently head of the Biochemical Assay Development and High Throughput Screening Unit, and am a Project Leader in the Oncology Department of Hoffmann-LaRoche, assignee of the above-captioned patent application.

2. I have extensive experience in molecular biology and immunology. One of my prime research interests has been the study of cytokines and their role in signal transduction-mediated regulation of normal and pathologic immune-related processes. In particular, I have been extensively involved in the study of the cytokine IL-12, which is also known as natural killer stimulatory factor (NKSF) and cytotoxic lymphocyte maturation factor (CLMF), and have published

10675 U.S. PTO
09/401839
09/22/99

numerous research articles relating to this topic. A copy of my curriculum vitae is attached hereto as Exhibit 1.

3. I hold a Ph.D. in Pharmacology, which I received in 1985 from Harvard University. From 1985 to 1988 I was a postdoctoral fellow in the laboratory of Dr. Ethan M. Shevach, Lab of Immunology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD.

4. I joined Hoffmann-LaRoche ("Roche") in 1988 as a senior scientist in Molecular Genetics. From 1991 to 1998 I was affiliated with Roche's Inflammation/Autoimmune Diseases division. In 1998 I began my current position as a project leader in Roche's Oncology Department, and in 1999 began my present, concurrent position as Head of Roche's Biochemical Assay Development and High Throughput Screening Unit.

5. As summarized on the attached curriculum vitae, I have authored or co-authored numerous scientific publications, have received several honors and awards and belong or serve on various professional societies and committees.

6. I have read and am familiar with United States Patent No. 5, 811,523 (the "523 patent"), which describes a purification and cloning of NKSF. As discussed below, with respect to the 523 patent's discussion of antibodies directed against NKSF, it is clear that the 523 patent neither

adequately teaches how to make the full scope of the antibodies claimed, nor does it teach a practical way of using antibodies which specifically bind NKSF. In particular, the '523 patent provides neither a practical therapeutic use nor a practical diagnostic use for such antibodies.

7. The '523 patent states that NKSF antibodies can be generated using "standard methods for diagnostic or therapeutic use" ('523 patent c. 10, 11. 25-28). No other discussion, however, relating to the use of NKSF antibodies as part of a therapeutic use is to be found within the '523 patent. The '523 patent only proposes that certain diseases or disorders (e.g., cancer, bacterial or viral infections, immune cell deficiencies) could be treated by administering NKSF polypeptides or fragments thereof.

8. Importantly, the therapeutic uses stated in the '523 patent require an increase in NKSF activity or an enhancement of natural killer (NK) cells, a cell type whose activity is increased by NKSF. Because of this, even if NKSF antibody administration had been contemplated by the '523 patent, NKSF antibodies, especially NKSF antibodies generated using "standard methods," could not be used to increase NKSF activity or enhance NK cell function. Rather, administration of NKSF antibodies would either have no effect on either NKSF activity or NK cell function, or, if an effect was observed, administration of NKSF antibodies would actually decrease or

abolish (i.e., "neutralize") NKSF activity or NK cell function.

9. In order for an NKSF antibody to increase NKSF activity and, therefore, enhance NK cell function, the antibody would be required to at least partially mimic or induce NKSF action. An antibody directed against a soluble factor such as NKSF, however, would, if anything, exert the opposite effect. For example, an antibody which specifically binds NKSF could act to inhibit or weaken binding of NKSF to the cellular receptor or receptors to which it would normally bind and through which it would normally act to enhance NK cell function.

10. The only antibody mentioned in the '523 patent is, in fact, reported to exhibit a negative effect on NKSF activity ('523 patent, c. 16, ll. 40-46). In particular, the '523 patent states that a polyclonal antiserum directed against the NKSF 40 kD subunit blocks NKSF activity in an in vitro assay. I note, however, that the NKSF-related specificity of the reported result is questionable. Specifically, the '523 patent presents no controls nor is characterization of the antiserum provided. The only indication that an antibody which specifically reacts with NKSF might be present in the antiserum is based on the antiserum's purported NKSF inhibitory activity. This is particularly true in view of the fact that a high antiserum concentration (only a 1:100 dilution) was utilized, which can often lead to artifacts, coupled with the fact that no

controls were presented which could have ruled out possible artifacts.

11. The '523 patent also fails to teach practical diagnostic uses employing NKSF antibodies. First, the '523 patent is absolutely silent regarding diagnostic uses whatsoever, except for the statement that diagnostic uses could be performed. The '523 patent provides no information, however, that would adequately teach one of skill in the art how an NKSF antibody could practically be used for a diagnostic purpose.

12. In particular, the '523 patent fails to teach such a diagnostic use could only be performed with certain types or sets of NKSF antibodies exhibiting particular characteristics. This is due to the fact that conditions that promote production of biologically active NKSF heterodimer also promote production and secretion of free NKSF 40 kD subunit, which does not exhibit the heterodimeric activity. Thus, a successful antibody-based diagnostic use would require employing an antibody or antibodies that can somehow distinguish between the the 70 kD NKSF heterodimer and the free 40 kD NKSF subunit (e.g., a single antibody or combination of antibodies which can distinguish between the 70 kD heterodimer and the free subunits, especially the 40 kD subunit). The '523 patent fails, however, to point out this requirement for a diagnostic use.

13. Finally, it is noted that the '523 patent does not provide an adequate level of teaching that would allow routine production of the full range of antibodies claimed. Taking just one example, the teaching provided in the '523 patent would not allow one of skill in the art to routinely produce antibodies, especially monoclonal antibodies, which specifically bind the 30-35 kD NKSF subunit. As mentioned above, the '523 patent merely states that NKSF polypeptides, in conjunction with standard methods, can be used to generate NKSF antibodies. For example, I do not know of a single instance where purified or recombinant heterodimeric NKSF has successfully been used to generate a monoclonal antibody against the 30-35 kD subunit of NSKF. The '523 patent fails to explicitly teach that would be a problem, and, further, also fails to explicitly teach alternative routes by which to generate monoclonal antibodies directed against this subunit, such as utilizing only the 30-35 kD subunit or a synthetic peptide thereof as a starting antigen.

14. I further declare that all statements made in this Declaration are of my own knowledge and are true and that all statements made on information and belief are believed to be true and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such

